

Influence of amino acid residues and their sequence in the molecules of dipeptides on the receptor properties and morphology of thin films based on them

© **Alsina A. Bismukhametova, Irina G. Efimova,¹⁺ Marat A. Ziganshin,^{1*}
Valeriy V. Gorbachuk,¹ Sufia A. Ziganshina,²
Anton P. Chuklanov,² and Anastas A. Bukharayev²**

¹ Department of Physical Chemistry. A.M. Butlerov Chemical Institute at KFU. Kremlevskaya St., 18.
Kazan, 420008. Tatarstan Republic. Russia. Phone: +7 (843) 233-73-09. Fax: +7 (843) 233-74-16.

² Laboratory of Physics and Chemistry of Surface. E.K. Zavoiskiy Kazan, KazSC RAS.
Sibirskiy trakt St., 10/7. Kazan, 420029. Tatarstan Republic. Russia. Phone: +7 (843) 231-91-07.

*Supervising author; +Corresponding author

Keywords: *Dipeptide, biozeolite, inclusion compounds, morphology of nanofilms, gravity sensors, atomic force microscopy.*

Abstract

By atomic force microscopy we studied the effect of methanol vapor on the morphology of thin films of the dipeptides of *L*-alanyl-*L*-valine, *L*-valyl-*L*-alanine and *L*-valyl-*L*-valine. Terms of saturation and adsorption capacity of dipeptides in relation to methanol have been determined by means of quartz microbalance.