

## Synthesis and structure of $\mu$ -oxobis[(4-iodphenoxy)tri-*n*-tolylantimony] [(4-MeC<sub>6</sub>H<sub>4</sub>)<sub>3</sub>SbOC<sub>6</sub>H<sub>4</sub>I-4)]<sub>2</sub>O

© Vladimir V. Sharutin,\*<sup>†</sup> Olga K. Sharutina,  
Antonina P. Pakusina, and Svetlana A. Smirnova

Department of Chemistry. Blagoveshchensk State Pedagogical University. Lenin St., 104.  
Blagoveshchensk, 675000. Russia. Phone: +7 (4162) 37-61-91. E-mail: vvsharutin@rambler.ru

\*Supervising author; <sup>†</sup>Corresponding author

**Keywords:**  $\mu$ -oxobis[(4-iodphenoxy)tri-*p*-tolylantimony], synthesis, structure.

### Abstract

Oxidation of tri(*p*-tolyl)antimony with hydrogen peroxide in ether in the presence of 4-iodophenol (1:1:1 mol.) is obtained  $\mu$ -oxobis[(4-iodphenoxy)tri-*p*-tolylantimony], in a centrosymmetric linear molecule which Sb atoms have a distorted trigonal-bipyramidal coordination with aryl ligands in equatorial positions, the bridging oxygen atom and aroxoligand in axial positions. Bond lengths of Sb-C, Sb-O<sub>bridge</sub>, Sb-O<sub>arox</sub> and angle SbOSb equal to 2.097(7)-2.114(7) $\text{\AA}$ ; 1.9978(6), 2.126(5) $\text{\AA}$  and 180°.