

Synthesis of new calyx[4]resorcines with sterically hindered phenol fragments

© Tatiana A. Barsukova,¹⁺ Sergey V. Bukharov,^{2*} Alexander R. Burilov,³
Gulnara N. Nugmanova,¹ Victor V. Syakaev⁴ and Natalia A. Mukmeneva¹

¹ Department of Synthetic Rubber Technology. Kazan National Research Technological University.
K. Marx St., 68. Kazan, 420015. Tatarstan Republic. Russia. Phone: +7 (843) 231-42-49.

E-mail: tanibars@yandex.ru

² Department of General Organic and Petrochemical Synthesis Technology. Kazan National Research Technological University. K. Marx St., 68. Kazan, 420015. Tatarstan Republic. Russia.

Phone: +7 (843) 231-45-78. E-mail: bukharov_@rambler.ru

³ Organoelemental Chemistry Division. A.E. Arbuzov Institute of Organic and Physical Chemistry Kazan Scientific Center of the Russian Academy of Science. Akad. Arbuzov St., 8. Kazan, 420088.

Tatarstan Republic. Russia. Phone: +7 (843) 272-73-44. E-mail: burilov@iopc.ru.

⁴ Physico-Chemical Research Division. A.E. Arbuzov Institute of Organic and Physical Chemistry Kazan Scientific Center of the Russian Academy of Science. Akad. Arbuzov St., 8. Kazan, 420088.

Tatarstan Republic. Russia. Phone: +7 (843) 262-68-21. E-mail: vsyakaev@iopc.ru.

*Supervising author; ⁺Corresponding author

Keywords: sterically hindered phenols, calyx[4]resorcines, synthesis, ¹H NMR.

Abstract

New calyx[4]resorcines with sterically hindered phenol fragments on the lower rim were synthesized by condensing resorcinol and methyl ester of resorcinol with sterically hindered benzaldehydes. Elucidation of their structures was made by NMR ¹H spectroscopy, Maldi Tof, IR spectra.