

## Synthesis of colloid silsesquioxanes based on organosilicon derivatives of *p*-tert-butyl thiacalix[4]arene

© Vladimir V. Gorbachuk,<sup>1</sup> Lyudmila S. Yakimova,<sup>1</sup> Alena A. Vavilova,<sup>1</sup>  
Ramilia V. Ziatdinova,<sup>1</sup> Ildar Ha. Rizvanov,<sup>2\*†</sup> and Ivan I. Stoikov<sup>1,3\*†</sup>

<sup>1</sup>Department of Organic Chemistry. A.M. Butlerov Chemical Institute. Kazan (Volga Region) Federal University. Kremlevskaya St., 18. Kazan, 420008. Russia. Phone: +7 (843) 233-74-62.

E-mail: [ivan.stoikov@mail.ru](mailto:ivan.stoikov@mail.ru)

<sup>2</sup>Arbuzov Institute of Organic and Physical Chemistry of Kazan Scientific Center of RAS.

Ac. Arbuzov St., 8. Kazan, 420088. Russia.

<sup>3</sup>Kazan institute of biochemistry and biophysics of Kazan Scientific Center of RAS.

Lobachevskogo St., 2/31. Kazan, 420111. Russia.

\*Supervising author; †Corresponding author

**Keywords:** thiacalixarene, colloid particles, silsesquioxanes, dynamic light scattering.

### Abstract

We have developed a method allowing to attain monodisperse colloid systems of submicron and nanosized particles based on tetrasubstituted organosilicon derivatives of *p*-tert-butyl thiacalix[4]arene in *cone*, *partial cone* and *1,3-alternate* conformations. Colloid systems were characterized by <sup>1</sup>H NMR-spectroscopy and dynamic light scattering method.