Thema	itic .	Section:	Prepar	rative	Che	mistr	y.					Full Paper
G 1		_	1 -	. 1 .	~1	. , '	-	 <u> </u>	 	 1	CD 11:	 12 22 2 5

Subsection: Organoelemental Chemistry.

Registration Code of Publication: 13-33-2-55

Publication is available for discussion in the framework of the on-line Internet conference "New methods of

Publication is available for discussion in the framework of the on-line Internet conference "New methods of synthesis, structure and application of organoelemetal compounds" http://butlerov.com/synthesys/

Contributed: February 7, 2013.

Synthesis and structure of μ -oxo[tris(2-methoxy,5-bromophenyl)trifluoroacetatoantimony]

© Vladimir V. Sharutin,* Vladislav S. Senchurin, and Olga K. Sharutina

Department of Organic Chemistry. Chemistry Faculty. Research South Ural State University. V.I. Lenin St., 76. Chelyabinsk, 454080. Russia. Phone: +7 (351) 267-95-70. E-mail: vvsharutin@rambler.ru

*Supervising author; *Corresponding author

Keywords: μ -oxo[tris(2-methoxy,5-bromophenyl)trifluoroacetatoantimony], synthesis, structure.

Abstract

 μ -Oxo[tris(2-methoxy,5-bromophenyl)trifluoroacetatoantimony] is obtained in 92% yield from equimolar amounts of tris(2-methoxy,5-bromophenyl)antimony, trifluoroacetic acid and hydrogen peroxide. According to preliminary data of X-ray analysis antimony atoms in molecules have a trigonal-bipyramidal coordination with the oxygen atoms in axial positions. Bond lengths of Sb-C, Sb-O_{bridge}, Sb-O_{term} and the angle SbOSb are 2.106(10)-2.139(11); 1.919(10), 1.932(10); 2.135(12)-2.334(12) Å and 168.05(8)° respectively.