

Synthesis and molecular structure of 4-acetyl-5-carboxymethyl-5-methyl-2-oxo-3-phenyloxolane

© Evgenia A. Zinina,¹ Vitaly V. Sorokin,¹ and Zoya A. Starikova²

¹ Department of Organic and Bioorganic Chemistry. N.G. Chernyshevsky Saratov State University.
Astrakhanskaya St., 83. Saratov, 410012. Russia. Phone: +7 (8452) 27-14-91.

E-mail: sorokinvv@info.sgu.ru, evgeniyaz@rambler.ru

² Laboratory of X-ray Diffraction. A.N. Nesmeyanov Institute of Organoelemental Compounds
RAS. Vavilov St., 28, B-334. Moscow, 119991. Russia.

Phone: +7 (499) 135-92-71. E-mail: star@xray.ineos.ac.ru

*Supervising author; †Corresponding author

Keywords: 4-acetyl-5-carboxymethyl-5-methyl-2-oxo-3-phenyloxolane, synthesis, X-ray structure analysis.

Abstract

4*t*-Acetyl-5*c*-carboxymethyl-5*t*-methyl-2*t*-oxo-3*r*-phenyloxolane with oxolane ring in the form of "half-chair", *trans*-positioned pseudoequatorial phenyl – and acetyl group was obtained by peroxide oxidation of 2*t*,4*t*-diacetyl-5*t*-hydroxy-5*c*-methyl-3*r*-phenylcyclohexanone.