Thematic Section: Preparative Research	·	Full Paper

Subsection: Organoelemental Chemistry.

Registration Code of Publication: 13-36-11-83

Publication is available for discussion within the functioning of the permanent internet-Conference "New methods of synthesis, structure and application of organoelemental compounds" http://butlerov.com/synthesys/Contributed: October 26, 2013.

Synthesis and structure of μ -oxo-bis-[(trifluoroacetato)-(tri-m-tolyl)antimony][(C₆H₄Me-3)₃SbOC(O)CF₃)]₂O

© Vladimir V. Sharutin,* Olga K. Sharutina, and Vladislav S. Senchurin
Department of Organic Chemistry. Chemistry Faculty. South Ural State University. Pr. Lenin, 76.
Chelvabinsk, 454080. Russia. Phone: +7 (351) 267-95-70. E-mail: vvsharutin@rambler.ru

*Supervising author; *Corresponding author

Keywords: synthesis, structure, [(C6H4Me-3)3SbOC(O)CF3)]2O.

Abstract

By reaction of (tri-*m*-tolyl)antimony with trifluoroacetic acid in the presence of hydrogen peroxide in aqueous-ether solution we obtained μ -oxo-bis-[(trifluoroacetate)(tri-*m*-tolyl)antimony] (**I**) with the yield of 91%. In the dinuclear molecule **I** SbOSb angle is 146.0(5)°. Antimony atoms have a distorted trigonal-bipyramidal coordination. Distances Sb-C vary in the range 2.086(7)-2.118(7) Å. Bonds of the antimony atom with bridging oxygen atom [1.962(6), 1.957(6) Å] are shorter than with oxygen atoms of trifluoroacetate groups [2.209(5), 2.209(5) Å].