

Bicyclic anionic σ -adduct of 2-hydroxy-3,5-dinitropyridine in the synthesis of new derivatives of 2,6-diazatricyclododekanes

© Evgenia V. Ivanova,^{1†} Irina I. Surova,¹ Nikolay V. Hlytin,¹ Igor V. Blokhin,¹ Irina V. Shakheldyan,¹ Yuriy M. Atroshchenko,^{1*†} and Konstantin I. Kobrakov^{2*}

¹Department of Chemistry. Tula State Lev Tolstoy Pedagogical University.

Lenin pr., 125. Tula, 300026. Russia. Phone: +7 (4872) 35-78-08. E-mail: reaktiv@tspu.tula.ru

²Department of Organic Chemistry. Moscow State University of Design and Technology.

Sadovnicheskaya St., 33. Moscow, 117997. Russia. Phone: +7 (495) 955-35-58. E-mail: kobrakovk@mail.ru

*Supervising author; †Corresponding author

Keywords: anionic σ -adducts, complex Yanovsky, 2,6-diazatricyclododekanes, 2-hydroxy-3,5-dinitropyridine.

Abstract

Studied the chemical behavior of the 2-hydroxy-3,5-dinitropyridine in reaction with carbanion of acetone in the Yanovsky reaction conditions, identified structure thus formed bicyclic anionic complex by IR and NMR-spectroscopy has been studied.