

## Phase equilibria in condensed systems with cyclododecane and *n*-alkanes

© Ivan K. Garkushin,<sup>1\*</sup> Alexander V. Kolyado,<sup>2+</sup> Evgeny P. Petrov,<sup>3+</sup>  
and Alexander A. Shamitov<sup>4+</sup>

Department of General and Inorganic Chemistry. Samara State Technical University. Molodogvardeiskaya  
St., 244, Samara, 443100. Samara Region. Samara. <sup>1)</sup> Phone: +7 (846) 278-44-77.

E-mail: <sup>1)</sup>baschem@samgtu.ru; <sup>2)</sup>kolyado@rambler.ru; <sup>3)</sup>e.p.petrov@yandex.ru; <sup>4)</sup>sansher@mail.ru;

\*Supervising author; <sup>+</sup>Corresponding author

**Keywords:** *n*-decane, *n*-octadecane, cyclododecane, phase equilibria, liquidus, eutectics.

### Abstract

By low-temperature differential thermal analysis using a differential scanning calorimeter of heat flow the systems of *n*-decane–*n*-octadecane–cyclododecane, *n*-decane–cyclododecane, *n*-octadecane–cyclododecane have been investigated. All of the studied systems belong to the eutectic type. The eutectic composition alloys contain 10.0 % wt in the system *n*-decane–*n*-octadecane–cyclododecane; eutectic composition alloy contains 84.5 % wt. *n*-C<sub>10</sub>H<sub>22</sub>; 5.5 % wt. *n*-C<sub>18</sub>H<sub>38</sub>; 10 % wt. C<sub>12</sub>H<sub>24</sub> and has the melting point of 34.9 °C.