

Synthesis of new thiazoles based on the reaction of *N*-aryl-*N'*-(quinazoline-2-yl) amidinothioureas with α -halocarbonyl compounds

© Khidmet S. Shikhaliev,* Alexey A. Kruzhilin,*[†]

Nadezhda V. Stolpovskaya,* and Alexander V. Falaleev

Organic Chemistry Department. Voronezh State University. University Square, 1.

Voronezh, 394018. Russia. Phone.: +7 (4732) 20-84-33. E-mail: chocd261@chem.vsu.ru

*Supervising author; [†]Corresponding author

Keywords: hetaryl amidinothiourea, heterocyclization, thiazoles.

Abstract

Biological activity of heterocyclic systems containing quinazoline and thiazole cycles widely studied. In this regard the synthesis of polyheterocyclic compounds with those cycles is an important problem to research novel biologically active substances. The purpose of this work is the synthesis of new quinazolyl-2-yl amidinothioureas and study of their reactions with α -haloketones: phenacyl bromide, ethyl monochloroacetate, 2-chloroacetoacetate and 3-chloroacetylacetone. Structures of the obtained compounds are confirmed by ¹H NMR spectrometry and elemental analysis. Probability of obtaining a bioactivity to inhibit 2-gluconate dehydrogenase virtually determined.

A two-step synthesis of heterocyclic systems with thiazole and quinazoline rings based on interaction of substituted amidinothioureas with α -halocarbonyl compounds such as phenacyl bromide, ethyl monochloroacetate, 3-chloroacetylacetone 2-chloroacetoacetate was studied. The optimal conditions for the reactions investigated: synthesis starting *N*-aryl-*N'*-quinazolyl 2-yl-thioureas was carried out at continuous stirring at room temperature in dimethylacetamide. Optimal conditions of interaction of α -halocarbonyl compounds with *N*-aryl-*N'*-quinazolylamidinothioureas is boiling in alcoholic medium followed by treatment with an aqueous alkali solution.

It is found that the reaction with phenacyl bromide, ethyl monochloroacetate, 3-chloroacetylacetone selectively proceed with thioamide moiety of quinazolyl-2-yl amidinothioureas. The first step of this process is the *S*-alkylation according to the mechanism described for the first time by Hantzsch. In the second step proceeds thiazole ring closure by nucleophilic addition with amino-group of formed intermediate. Nucleophilicity of the nitrogen atom linked to the aromatic system reduced due to the bias of electron density of the benzene ring, that's why secondary amino-group of guanidine moiety involved in the cyclization. Interaction of 2-quinazolylamidinothioureas with 2-chloroacetoacetate occurs not so definitely and may lead to the formation of two alternative products. It is shown that heterocyclization with ethyl-2-chloroacetoacetate affects carbonyl group and result in formation of ethyl 2-[(*R*₁-phenyl)imino]-3-{imino[(6-*R*₂-4-methyl-quinazolin-2-yl)-amino]methyl}-2,3-dihydro-1,3-thiazole-5-carboxylates.

References

- [1] R.C. Schnur, A.F.J. Fliri, S. Kajiji, V.A. Pollack. *N*-(5-fluorobenzothiazol-2-yl)-2-guanidinothiazole-4-carboxamide. A Novel, Systemically Active Antitumor Agent Effective against 3LL Lewis Lung Carcinoma. *J. Med. Chem.* **1991**. Vol.34. P.914-918.
- [2] Y. Katsura, Y. Inoue, T. Tomishi, H. Ishikawa, H. Takasugi. 2-Guanidino-4-pyridylthiazoles as Histamine H₂-Receptor Antagonists with Potent Gastroprotective Effects against Nonsteroidal Antiinflammatory Drug-Induced Injury. *J. Med. Chem.* **1994**. Vol.37. No.1. P.57-62.
- [3] V.K. Pandey, D. Misra, M.N. Joshi, K. Chandra. Quinazolyl benzophenothiazines as potential antiviral agents. *Pharmacological Research Communications.* **1988**. Vol.20. No.2. P.153-165.
- [4] K.M. Murav'eva, N.V. Arkhangel'skaya, M.N. Shchukina. Synthesis and tuberculostatic activity of *N*-phenyl-*N'*-(quinazolyl-6)-thioureas. *Pharmaceutical Chemistry Journal.* **1968**. Vol.2. No.1. P.1-3.
- [5] A.K. Pandey, R. Singh, V.K. Verma. Synthesis of some 3,5-diarylimino-2-phenyl-1,2,4-thiadiazolidines. Oxidative debenzylsulfon and cyclization of 1 (*N,N*-diarylamidino)-3-aryl-2benzylthioureas. *Acta Chim. Hung.* **1983**. Vol.114. No.2. P.107-111.

- SYNTHESIS OF NEW THIAZOLES BASED ON THE REACTION OF N-ARYL-N'-(QUINAZOLINE-2-YL)...* ____ 64-70
- [6] Y. Kihara, S. Kabashima, K. Uno. Oxidative heterocyclization using diethyl azodicarboxylate. *Synthesis*. **1990**. No.1. P.1020-1023.
- [7] G.R. Murthy, V.M. Reddy. New hypoglycemic agents. Part III. Synthesis and hypoglycemic activity of some new 1-(2-benzimidazolamidino-3-aryl/cyclohexyl-2-thioureas. *Indian J. Pharm. Sci.* **1987**. Vol.49. No.5. P.175-177.
- [8] R.A. Cutler, S. Schalit. Amidinoureas and amidinothioureas: *Pat.* US3798269. IPC C07D295/13; (IPC1-7): C07C127/16. The applicant and patent holder Cutler R.A., Schalit S. No.USD3798269. appl.08.10.1970; published 19.03.1974. 14p.
- [9] H. Beyer, H. Hantschel. Über Thiazole, XL. Synthesen von Thiazolyl-2-guanidinen. *European Journal of Inorganic Chemistry*. **1962**. Vol.95. No.4. P.893-901.
- [10] H. Beyer, H. Hantschel. Über Thiazole, XLI. Thiazolyl-2-pyrimidyl-2-amine. *Chemische Berichte*. **1962**. Vol.95. No.4. P.902-906.
- [11] A. Hantzsch, J.H. Weber. Über Verbindungen des Thiazols (Pyridins der Thiophenreihe). *European Journal of Inorganic Chemistry*. **1887**. Vol.20. No.2. P.3118-3132.
- [12] J.V. Metzger. Thiazole and Its Derivatives: Part One. *John Wiley*. **1979**. 233p.
- [13] R.F. Hunter, E.R. Parken. The unsaturation and tautomeric mobility of heterocyclic compounds. Part IV. The methylation and bromination of a series of 2-*p*-substituted anilinothiazoles. *Journal of the Chemical Society (Resumed)*. **1934**. P.1175-1177.