

Indicative indicators of eutrophication as an element of monitoring of aquatic ecosystems

© Marina L. Kalayda,*⁺ and Madina F Khamitova

Department "Water Bioresources and Aquaculture". Kazan State Power Engineering University.

Krasnenskiy St., 51. Kazan, 420066. Republic of Tatarstan. Russia.

Phone: +7 (843) 519-43-53. E-mail: kalayda@mi.ru

*Supervising author; ⁺Corresponding author

Keywords: environmental pollution, eutrophication, Kuibyshev reservoir, wastewater, pollution indication, benthos.

Abstract

The choice of criteria for assessing the state of aquatic ecosystems has long been an important aspect of environmental and hydrobiological research. Changes in the mean multiyear values of the indices of chemical and biological oxygen consumption, characterizing the conditions of the degree of eutrophication, been shown. Using the ratio of MPC for COD and BOD5 defined "normative" ratio – 13.33%. It was shown that this ratio in control areas was 7.94-7.89%, in the zone of wastewater mixing with the water of the reservoir the ratio was 14.97, in the area of wastewater discharge from the primary settler this ratio was the maximum – 58.48%.

Since the level of phytoplankton development reflects the level of natural eutrophication of aquatic ecosystems, a comparative analysis of water quality indicators for the development of phytoplankton was conducted. It is shown that the calculated indices do not reflect the real ecological specifics of the sites.

Based on the study of the components of the hydrobiocenosis of the section of the Volga Reach in the area of wastewater discharges of the Mari Pulp and Paper Mill, it was shown that under the conditions of anthropogenic eutrophication, there is a significant variability in the characteristics of the hydrobiocenosis including species diversity. The coefficient of variation of the indices of species diversity better reflects the level of eutrophication.

A developed Index Y is proposed that reflects the ratio of the main complex of zoobenthos species with different types of respiration: in areas of the greatest anthropogenic eutrophication, it is represented mainly by insect species capable of atmospheric respiration; when the state in the ecotope is improved, the diversity and representation of the secondary-water insects increases, and the groups consuming oxygen dissolved in water replace forms that consume atmospheric air; primary-water animals predominate in the control areas and in the restoration zones. The values of the indicator can vary from 100 if there are no insects in the zoobenthos to 0 in the absence of zoobenthos or if it is represented exclusively by hydrobionts, which are adapted to breathing atmospheric air. The main criterion for the state of the system is the ratio of zoobenthos to oxygen in water, therefore this index can be considered as an indicator of eutrophication of the ecosystem. The lower the index, the more critical the state of the aquatic ecosystem. The index, unlike the others, is not an indicator of the ongoing process of eutrophication, but reflects the already negative impact on the aquatic ecosystem and assesses its ability to restore.

References

- [1] M.L. Kalayda. Ensuring water quality in the Republic of Tatarstan is the global civil duty of every modern person. The Earth Charter is a practical tool for solving fundamental problems of sustainable development: a collection of materials of the International Scientific and Practical Conference dedicated to the 15th anniversary of the implementation of the principles of the Earth Charter in the Republic of Tatarstan. *Kazan: Tatar. Book publ. house. 2016*. P.148-152. (russian)
- [2] M.L. Kalayda. Aquaculture as a basis for improving water quality in the Republic of Tatarstan. *Newsletter SC SoL. 2016*. No.3(29). P.115-122. (russian)
- [3] M.L. Kalayda, Al-Bachri Walid Sami Jawad. Regulation and quality control of waters as an important component of ensuring public health. Environmental problems and public health a collection of articles of the All-Russian Scientific and Practical Conference. *Penza: ED PGACA. 2016*. P.34-38. (russian)

- [4] M.L. Kalayda, M.F. Khamitova. Possibilities of applying eichornia in the post-treatment of waters of the pulp and paper mill. Part 1. Features of the chemical composition of water in the Volga reach of the Kuibyshev reservoir. *Butlerov Communications*. **2015**. Vol.44. No.11. P.97-103. DOI: 10.37952/ROI-jbc-01/15-44-11-97
- [5] M.L. Kalayda, M.F. Khamitova. Possibilities of applying eichornia in the post-treatment of waters of the pulp and paper mill. Part 2. Peculiarities of chemical composition of bottom sediments in the Volga reach of the Kuibyshev Reservoir. *Butlerov Communications*. **2015**. Vol.44. No.11. P.104-112. DOI: 10.37952/ROI-jbc-01/15-44-11-104
- [6] M.L. Kalayda, M.F. Khamitova. Possibilities of applying eichornia in the post-treatment of waters of the pulp and paper mill. Part 3. Features of the chemical composition of *Eichornia crassipes*. *Butlerov Communications*. **2015**. Vol.44. No.11. P.113-121. DOI: 10.37952/ROI-jbc-01/15-44-11-113
- [7] M.L. Kalayda, M.F. Khamitova, S.I. Novotochinov. Change in the chemical index of oxygen during algolization of water by chlorella. *Butlerov Communications*. **2016**. Vol.48. No.10. P.143-149. DOI: 10.37952/ROI-jbc-01/16-48-10-143
- [8] V.K. Shitikov, G.S. Rosenberg, T.D. Zinchenko. Quantitative hydroecology: methods of systemic identification. *Tolyatti: IEVB RAS*. **2003**. 463p. (russian)