

The structure and stability of higher fullerenes C₁₀₄: IPR isomers 812 (D₂) and 822 (D_{3d})

© **Alsu A. Gaynullina**,¹ **Ayrat R. Khamatgalimov**,^{1,2} and **Valery I. Kovalenko***⁺

¹ Department of Industrial Ecology. Engineering Chemical & Technological Institute. KNRTU.

Karl Marx St., 68. Kazan, 420015. Tatarstan Republic. Russia.

Phone: +7 (843) 273-22-83. E-mail: m.mitroshkina@mail.ru.

² Laboratory of PhysicoChemical Analysis. A.E. Arbuzov Institute of Organic and Physical Chemistry.

Akad. Arbuzov St., 8. Kazan, 420088. Tatarstan Republic. Russia.

Phone: +7 (843) 273-22-83. E-mail: koval@iopc.ru

*Supervising author; ⁺Corresponding author

Keywords: fullerene C₁₀₄, isolated pentagon rule, molecular structure, bonds distribution, substructures, quantum-chemical calculations.

Abstract

An analysis of the molecular structures of two IPR isomers 812 (D₂) and 822 (D_{3d}) of higher fullerene C₁₀₄ is carrying out. The data about the distribution of single bonds, double bonds, and delocalized in hexagon π -bonds in studied isomer molecules are presented for the first time as well as their structural formulas instead of conventional molecular graphs. Quantum chemical calculations (DFT) show that isomer 812 (D₂) have a closed electronic shell, whereas isomer 822 (D_{3d}) contains a pair of equivalent radical substructures, each one consists of three condensed phenalenyls. The calculated energy difference between the two molecules is ca. 25 kcal/mol in favor of 812 (D₂) isomer. It is found that double bonds connecting pentagons in pristine isomer 812 (D₂) are most short ones similarly to previously studied higher fullerenes which is important for understanding the synthesis process of fullerene derivatives. The preliminary bonds distributions according to the developed approach are confirmed in general by DFT calculations. Some differences between calculated bond lengths and predicted ones should be explained in terms of great fullerene size and, consequently, great number of condensed hexagons (compared with constant number of pentagons, which are equal to 12). These bond lengths deviations are belonging to fragments of molecules that similar to nanotubes without pentagons. Our previous analysis of the addend distribution in a series of higher IPR fullerenes up to C₉₀ showed the additions were essentially to hexagons with delocalized π -bonds. We suppose that a similar picture of addends distribution would be in exohedral derivatives of higher fullerene C₁₀₄. The important practical result is that the disclosed structures of fullerene molecules open a conscious way of chemical modification of fullerene molecules to obtain important derivatives.

References

- [1] P.W. Fowler, D.E. Manolopoulos. An Atlas of Fullerenes. *Dover Publ., Mineola N.Y.* **2006.** 392p.
- [2] S. Yang, S. Wang, S.I. Troyanov. The most stable isomers of giant fullerenes C₁₀₂ and C₁₀₄ captured as chlorides, C₁₀₂(603)Cl_{18/20} and C₁₀₄(234)Cl_{16/18/20/22}. *Chem. Eur. J.* **2014.** Vol.20. P.6875-6878.
- [3] S. Yang, T. Wei, E. Kemnitz, S.I. Troyanov. First isomers of pristine C₁₀₄ fullerene structurally confirmed as chlorides, C₁₀₄(258)Cl₁₆ and C₁₀₄(812)Cl₂₄. *Chem. Asian J.* **2014.** Vol.9. P.79-82.
- [4] S. Wang, S. Yang, E. Kemnitz, S.I. Troyanov. New giant fullerenes identified as chloro derivatives: Isolated Pentagon-Rule C₁₀₈(1771)Cl₁₂ and C₁₀₆(1155)Cl₂₄ as well as nonclassical C₁₀₄Cl₂₄. *Inorg. Chem.* **2016.** Vol.55. No.12. P.5741-5743.
- [5] B.Q. Mercado, A. Jiang, H. Yang, Z. Wang, H. Jin, Z. Liu, M.M. Olmstead, A.L. Balch. Isolation and structural characterization of the molecular nanocapsule Sm₂@D_{3d}(822)-C₁₀₄. *Angew. Chem.Chem. Int. Ed.* **2009.** Vol.48. P.9114-9116.
- [6] N. Shao, Y. Gao, S. Yoo, W. An, X.C. Zeng. Search for lowest-energy fullerenes: C₉₈ to C₁₁₀. *J. Phys. Chem. A.* **2006.** Vol.110. P.7672-7676.

- [7] V.I. Kovalenko, M.V. Semyashova. Hexagon π -bond delocalization in fullerenes. *Chem. & Comput. Simulation. Butlerov Comm.* **2000**. No.3. P.41-43. (russian)
- [8] A.R. Khamatgalimov, V.I. Kovalenko. Characteristic reasons of fullerene instabilities on C_{72} and C_{74} . *Structure and dynamic of molecular system. Kazan, UNIPRESS.* **2003**. Vol.3. P.186-189. (russian)
- [9] V.I. Kovalenko, A.R. Khamatgalimov. Open-shell fullerene C_{74} : phenalenyl-radical substructures. *Chem. Phys. Lett.* **2003**. Vol.377. No.3-4. P.263-268. (russian)
- [10] V.I. Kovalenko, A.R. Khamatgalimov. Regularities in the molecular structures of stable fullerenes. *Russ. Chem. Rev.* **2006**. Vol.75. No.11. P.981- 988. (russian)
- [11] A.R. Khamatgalimov, V.I. Kovalenko. Molecular structures of unstable isolated-pentagon-rule fullerenes C_{72} - C_{86} . *Russ. Chem. Rev.* **2016**. Vol.85. No.8. P.836-853. (russian)
- [12] A. Rodriguez-Forteza, N. Alegret, A.L. Balch, J.M. Poblet. The maximum pentagon separation rule provides a guideline for the structures of endohedral metallofullerenes. *Nature Chem.* **2010**. Vol.2. P.955-961. (russian)
- [13] A. Streitwieser. Molecular orbital theory for organic chemists. *New York: Wiley.* **1961**. 489p.
- [14] D. Bochvar, E. Galpern. On hypothetical systems: carbon dodecahedron, S-icosahedron and carbon-S-icosahedron. *Proceedings USSR Acad. Sci.* **1973**. Vol.209. No.3. P.610-612. (russian)
- [15] R. Taylor. C_{60} , C_{70} , C_{78} , and C_{84} : numbering, π -bond order calculations and addition pattern considerations. *Chem. Soc. Perkin Trans.* **1993**. Vol.2. P.813-824.
- [16] V.I. Kovalenko, A.R. Khamatgalimov, A.G. Shamov. Electron-deficiency local substructures and stability of endohedral metallofullerenes of C_{82} . *Structure and dynamics of molecular system. Ioshkar-Ola: MarPTI Publ.* **2001**. Vol.8 (1). P.254-258. (russian)
- [17] V.I. Kovalenko. Comment on P. Bultinck's paper "Critical analysis of the local aromaticity concept in polyaromatic hydrocarbons". *Faraday Discuss.* **2007**. Vol.135. P.392-393.
- [18] A.A. Goryunkov. *CageGen2*, v. 0.015, **2015**. aag@thermo.chem.msu.ru
- [19] A.D. Becke. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**. Vol.98. P.5648-5652.
- [20] C. Lee, W. Yang, R.G. Parr. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys. Rev. B.* **1988**. Vol.37. P.785-789.
- [21] Gaussian 03 / M.J. Frisch, G.W. Trucks [et. al]; Gaussian, Inc. Pittsburgh PA. **2003**.
- [22] D. Tomanek. Guide through the nanocarbon jungle. *San Rafael, CA: Morgan & Claypool Publ.* **2014**. 93p.
- [23] L.H. Ye, B.G. Liu, D.S. Wang. Ab initio molecular dynamics study on small carbon nanotubes. *Chin. Phys. Lett.* **2001**. Vol.18. No.11. P.1496-1499.
- [24] A.N. Imtani, V.K. Jindal. Characterizing single-walled carbon nanotubes by pressure probe. *Carbon.* **2009**. Vol.47. P.3247-3251.
- [25] M.V. Mitroshkina, A.R. Khamatgalimov, V.I. Kovalenko. Molecular structure of the two IPR isomers 234 (C_s) and 258 (C_1) of a higher fullerene C_{104} . *Butlerov Communications.* **2017**. Vol.49. No.1. P.28-33. DOI: 10.37952/ROI-jbc-01/17-49-1-28
- [26] R.A. Tuktamysheva, A.R. Khamatgalimov, V.I. Kovalenko. Electronic and geometric structure of a number of fullerene isomers C_{90} and structure of their chlorine and perfluoroalkyl polyadducts. *Butlerov Communications.* **2014**. Vol.37. No.1. P.1-12. ROI: jbc-02/14-37-1-1