

Theoretical modeling of the interaction of 2-*R*-5,7-dinitrobenzo[d]-oxazoles with methoxide ion by DFT method

© Igor V. Blokhin,¹⁺ Loyk G. Mukhtorov,¹ Yury M. Atroshchenko,¹ Irina V. Shakhkeldyan,¹ Pavel V. Strashnov,² Mikhail A. Ryabov,² Konstantin I. Kobrakov,^{3*} and Alexey N. Shumsky⁴

¹Department of Chemistry. Tula State Lev Tolstoy Pedagogical University.

Pr. Lenina, 126. Tula, 300026. Russian Federation. Phone: +7 (4872) 35-78-08. E-mail: reaktiv@tspu.tula.ru

²Department of General Chemistry. Russian University of Peoples' Friendship. Mikluho-Maklaja St., 6.

Moscow, 117198. Russian Federation. Phone: +7 (495) 955-08-60. E-mail: maryabov@mail.ru

³Department of Organic Chemistry. Moscow State University of Design and Technology.

Sadovnicheskaya St., 33. Moscow, 117997. Russia. Phone: +7 (495) 955-35-58. E-mail: kobrakovk@mail.ru

⁴Emanuel Institute of Biochemical Physics of Russian Academy of Sciences.

Kosygina St., 4. Moscow, 119334. Russia. Phone: +7 (495) 939-73-66. E-mail: shumsky@mail.ru

*Supervising author; [†]Corresponding author

Keywords: 5,7-dinitrobenzo[d]oxazole, 2-methyl-5,7-dinitrobenzo[d]oxazole, 2-phenyl-5,7-dinitrobenzo[d]oxazole, nucleophilic addition, anionic σ -adduct, density functional theory, DFT/B3LYP.

Abstract

The interaction of 5,7-dinitrobenzo[d]oxazole, 2-methyl-5,7-dinitrobenzo[d]oxazole and 2-phenyl-5,7-dinitrobenzo[d]oxazole with a methoxide ion has been studied. It has been established that the nucleophile is attached to the carbon atom at position 2 of the heteroaromatic ring, and not to positions 4 and 6 with the formation of classical Mesenheimer complexes. The structure of the obtained σ -adducts was established by IR and NMR spectroscopy methods, as well as by elemental analysis data. Quantum-chemical calculations by the DFT method show that the largest positive charges are concentrated on C2, C3a and C7a atoms in the gas phase and methanol, whereas the presence of a relatively high electron density is observed on carbon atoms C4 and C6. Charges in methanol are slightly higher than in the gas phase. These data do not agree with the results of the calculation by the semiempirical method PM6, obtained earlier. Thus, it is established that the studied reaction proceeds under charge control conditions. The calculation of the total energies of the putative products showed that among the products of addition of the methoxide ion in the gaseous phase to the unsubstituted substrate, the most stable adducts are formed upon attack of the nucleophile at position 2, whereas in the case of 2-phenyl-5,7-dinitrobenzo[d]oxazole, the lowest values of the energy are the anions formed upon the addition of the methoxide ion to position 4. For a substrate with a methyl group at position 2, the difference in the energies of the adducts when the nucleophile is added to positions 2 and 4 is insignificant and is of the order of 0.5 kJ/mol. The attack of the methoxide ion on the carbon atom C-4, leading to the formation of classical adducts of Mesenheimer, is most energetically favorable in methanol for all substrates. Analysis of the contribution of the C-4 carbon atom to the LUMO substrate indicates the possibility of an attack of the methoxide ion over the benzannelated ring.

References

- [1] S.-T. Huang, I.J. Hsei, C. Chen. Synthesis and anticancer evaluation of bis(benzimidazoles), bis(benzoxazoles), and benzothiazoles. *Bioorganic & Medicinal Chemistry*. **2006**. Vol.14. No.17. P.6106-6119.
- [2] D. Kumar, M.R. Jacob, M.B. Reynolds, S.M. Kerwin. Synthesis and evaluation of anticancer benzoxazoles and benzimidazoles related to UK-1. *Bioorganic & Medicinal Chemistry*. **2002**. Vol.10. No.12. P.3997-4004.

- [3] M.L. McKee, S.M. Kerwin. Synthesis, metal ion binding, and biological evaluation of new anticancer 2-(2'-hydroxyphenyl)benzoxazole analogs of UK-1. *Bioorganic & Medicinal Chemistry*. **2008**. Vol.16. No.4. P.1775-1783.
- [4] E. Oksuzoglu, B. Tekiner-Gulbas, S. Alper, O. Temiz-Arpaci, T. Ertan, I. Yildiz, N. Diril, E. Sener-Aki, I. Yalcin. Some benzoxazoles and benzimidazoles as DNA topoisomerase I and II inhibitors. *Journal of Enzyme Inhibition and Medicinal Chemistry*. **2008**. Vol.23. No.1. P.37-42.
- [5] M.A. Weidner-Wells, K.A. Ohemeng, V.N. Nguyen, S. Fraga-Spano, M.J. Macielag, H.M. Werblood, B.D. Foleno, G.C. Webb, J.F. Barrett, D.J. Hlasta. Amidino benzimidazole inhibitors of bacterial two-component systems. *Bioorganic & Medicinal Chemistry Letters*. **2001**. Vol.11. No.12. P.1545-1548.
- [6] M.H. Potashman, J. Bready, A. Coxon, T.M. DeMelfi, L. DiPietro, N. Doerr, D. Elbaum, J. Estrada, P. Gallant, J. Germain, Y. Gu, J.-C. Harmange, S.A. Kaufman, R. Kendall, J.L. Kim, G.N. Kumar, A.M. Long, S. Neervannan, V.F. Patel, A. Polverino, P. Rose, S. van der Plas, D. Whittington, R. Zanon, Zhao H. Design, Synthesis, and Evaluation of Orally Active Benzimidazoles and Benzoxazoles as Vascular Endothelial Growth Factor-2 Receptor Tyrosine Kinase Inhibitors. *Journal of Medicinal Chemistry*. **2007**. Vol.50. No.18. P.4351-4373.
- [7] T. Ogoshi, J. Miyake, Y. Chujo. Multiresponsive Photopatterning Organic-Inorganic Polymer Hybrids Using a Caged Photoluminescence Compound. *Macromolecules*. **2005**. Vol.38. No.10. P.4425-4431.
- [8] T. Ogura, K.-t. Yamaguchi, Y. Shibasaki, M. Ueda. Photosensitive Poly(benzoxazole) Based on Poly(o-hydroxy amide), Dissolution Inhibitor, Thermoacid Generator, and Photoacid Generator. *Polymer Journal*. **2007**. Vol.39. No.3. P.245-251.
- [9] S. Park, S. Kim, J. Seo, S.Y. Park. Strongly Fluorescent and Thermally Stable Functional Polybenzoxazole Film: Excited-State Intramolecular Proton Transfer and Chemically Amplified Photopatterning. *Macromolecules*. **2005**. Vol.38. No.11. P.4557-4559.
- [10] D. Joule, K. Mills. Chemistry of heterocyclic compounds. *Moscow: Mir*. **2004**. 365p. (russian)
- [11] H. Bredereck, R. Gompper, F. Reich, U. Gotsma. Ringaufspaltung von Oxazolen mit 2,4-Dinitrophenylhydrazin. *Chemische Berichte*. **1960**. Vol.93. No.9. P.2010-2015.
- [12] E.V. Ivanova, I.I. Surova, N.V. Hlytin, I.V. Blokhin, I.V. Shakheldyan, Yu.M. Atroshchenko, and K.I. Kobrakov. Bicyclic anionic σ -adduct of 2-hydroxy-3,5-dinitropyridine in the synthesis of new derivatives of 2,6-diazatricyclododekanes. *Butlerov Communications*. **2014**. Vol.38. No.6. P.64-68. ROI: jbc-02/14-38-6-64
- [13] I.I. Surova, E.V. Ivanova, I.V. Blokhin, I.V. Shakheldyan, Yu.M. Atroshchenko, K.I. Kobrakov, D.N. Kuznetsov, and I.V. Fedyanin. Synthesis 6-thiosubstituted 3,5-dinitro-1,2,3,4-tetrahydropyridines. *Butlerov Communications*. **2015**. Vol.42. No.4. P.91-95. ROI: jbc-02/15-42-4-91
- [14] L.G. Mukhtorov, I.V. Blokhin, I.V. Shakheldyan, Yu.M. Atroshchenko, V.A. Arlyapov, K.I. Kobrakov, and A.N. Shumsky. Quantum chemical and experimental study of anionic σ -adducts of 2-methyl-5,7-dinitrobenzo[d]oxazole with methoxide ion. *Butlerov Communications*. **2015**. Vol.44. No.12. P.164-169. ROI: jbc-02/15-44-12-164
- [15] A.D. Becke. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**. Vol.98. No.7. P.5648-5652.
- [16] C. Lee, W. Yang, R.G. Parr. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Physical review B*. **1988**. Vol.37. No.2. P.785.
- [17] P.J. Stephens, F.J. Devlin, C.F. Chabalowski, M.J. Frisch. Ab initio calculation of vibrational absorption and circular dichroism spectra using density functional force fields. *The Journal of Physical Chemistry*. **1994**. Vol.98. No.45. P.11623-11627.
- [18] A. Schäfer, C. Huber, R. Ahlrichs. Fully optimized contracted Gaussian basis sets of triple zeta valence quality for atoms Li to Kr. *The Journal of Chemical Physics*. **1994**. Vol.100. No.8. P.5829-5835.
- [19] E.D. Glendening, J.K. Badenhoop, A.E. Reed, J.E. Carpenter, J.A. Bohmann, C.M. Morales, F. Weinhold. NBO 5.G. **2004**. URL: <http://www.chem.wisc.edu/~nbo5>.
- [20] A.A. Granovsky. Firefly version 7.1.G. **2016**. URL: <http://classic.chem.msu.su/gran/firefly/index.html>.
- [21] I.A. Pearl, W.M. Dehn. Derivatives of Picramic Acid and Some of their Rearrangements. *JACS*. **1938**. Vol.60. No.4. P.925-927.
- [22] Yu.M. Atroshchenko, S.N. Nasonov, S.S. Gitis, A.Ya. Kaminsky, A.I. Melnikov, I.V. Shakheldian. On the interaction of 1,3,5-trinitrobenzene with sodium tetrahydride borate. *Journal of Organic Chemistry*. **1994**. Vol.30. No.4. P.632-633. (russian)