

Thermodynamic properties of binary and ternary mixtures containing cellulose, caffeine, carbon dioxide and water, in supercritical fluid conditions

© Zufar I. Zaripov,¹ Lenar Yu. Yarullin,¹⁺ Farid M. Gumerov,^{1*} Truong Nam Hung,² and Farizan R. Gabitov¹

¹ Department of Theoretical Foundations of Heat Engineering. Kazan National Research Technological University. Karl Marx St., 68. Kazan, 420015. Republic of Tatarstan. Russia.

Phone: +7 (843) 231-42-11. E-mail: yarul.lenar@gmail.com

² Hanoi.Electric Power University. Hanoi. Vietnam.

*Supervising author; +Corresponding author

Keywords: caffeine, heat capacity, enthalpy, supercritical carbon dioxide, cellulose, mixing enthalpy.

Abstract

In this paper, we have studied the literature data of thermophysical properties as individual components, such as caffeine and cellulose, as well as binary and ternary systems containing tea leaf components. Also, data on the heat capacity of pure caffeine of various authors.

On the automated experimental setup created on the basis of the scanning calorimeter, the heat capacity of pure caffeine was obtained in the temperature range 323-523K. The densities of binary and ternary systems are determined. Measurements of the heat capacity and the density of the binary mixture caffeine-supercritical carbon dioxide (SC-CO₂) system are performed at pre- and supercritical pressures. The calculation formula for determining the specific heat of the component. The dependences of the heat capacity of the caffeine-SC-CO₂ system at various temperatures and pressures are graphically shown. Also graphically shows the dependence of the density of the caffeine-SC-CO₂ system on the pressure and at temperatures of 308, 323 and 343 K.

The results of measurements of the heat capacity of a mixture of caffeine-SC-CO₂ are recalculated into an excess heat capacity and are shown in the graph in the form of the dependence of the excess heat capacity of the system on the pressure at temperatures 308.323 and 343 K. According to the obtained values of the excess heat capacity of the caffeine-SC-CO₂ system, the enthalpy of dissolution of the system at various pressures and temperatures. Comparison of calculated and experimental data of the enthalpy of dissolution of the caffeine-SC-CO₂ system is shown graphically and indicates qualitative agreement.

The results of measuring the enthalpy of mixing of SC-CO₂ in an aqueous caffeine solution at various temperatures are presented. Comparison of the enthalpy of dissolution of CK-CO₂ in caffeine, cellulose, tea leaf and caffeine aqueous solution at a temperature of 308 K.

The method of determining the excess enthalpy associated with the calculation from the data of solubility is shown in the work. On the basis of literature data on solubility, molar volume and vapor pressure, enthalpy values were calculated. The results of the calculation and their comparison with the experimental data are shown in graphs.

References

- [1] N.H. Truong, F. Gumerov, F. Gabitov, R. Usmanov, V. Hairutdinov, B. Le Neindre. Improvement of the water brewing of Vietnamese green tea by pretreatment with supercritical carbon dioxide. *J. of Supercritical Fluids*. **2012**. Vol.62. P.73-78.
- [2] G.I. Burgos-Solyrzano, J.F. Brennecke, M.A. Stadtherr. Solubility measurements and modeling of molecules of biological and pharmaceutical interest with supercritical CO₂. *Fluid Phase Equilibrium*. **2004**. Vol.220. P.55-67.
- [3] J.R. Kim, J.B. Kyong. Thermodynamic Properties of Caffeine in Compressed Gas. *Bul.Korean Chem.Soc.* **1995**. Vol.16. No.5. P.432.
- [4] V.I. Kovalenko. Crystalline cellulose: structure and hydrogen bonds. *Chemical Success*. **2010**. Vol.79. No.3. P.261-272. (russian)

- [5] V.F. Uriyash, V.N. Larina, N.Y. Kokurina, N.V. Novoselova. Thermochemical characteristics of cellulose and its aqueous mixtures. *Journal of Physical Chemistry*. **2010**. Vol.84. No.6. P.1023-1029. (russian)
- [6] M.Ya. Eoyelovich. Enthalpy of formation and dissolution of cellulose. Union conference on chemical thermodynamics and calorimetry (tesis). *GorkiyGGU*. **1988**. Part 2. P.189-191. (russian)
- [7] N.B. Vargaftik Handbook of thermophysical properties of gases and liquids. *Moscow: Science*. **1972**. 720p. (russian)
- [8] L. Dordain, J.-Y. Coxam, J.-P. Grolier. Measurement of isobaric heat capacities of gases from 323.15 to 573.15 K up to 30 MPa. *Rev. Sci. Instrum.* **1994**. Vol.65. No.10. P.3263-3267.
- [9] A. Cesaro, G. Staree. Thermodynamic properties of caffeine cristal forms. *J. Phys. Chem.* **1980**. Vol.84. P.1345-1346.
- [10] H. Bothe, H. K. Cammenga. Phase transitions and thermodynamic properties of anhydrous caffeine. *Journal of Thermal Analysis*. **1979**. Vol.16. P.267-275.
- [11] C.B. Atllio, Giorglo Starec. Thermodynamic properties of caffeine crystal forms. *J. Phys. Chem.* **1980**. No.84. P.1345-1346.
- [12] U.J. Griesser, M. Szelagiewicz, U.Ch. Hofmeier, C. Pitt and S. Cianferani. Vapor pressure and heat of sublimation of crystal polymorphs. *Journal of Thermal Analysis and Calorimetry*. **1999**. Vol.57. P.45-60.
- [13] G. Defossefont, L. Stanislaw. Randzio and B. Legendre. Contributions of calorimetry for Cp determination and of scanning transitiometry for the study of polymorphism. *Crystal Growth & Design*. **2004**. Vol.4. No.6. P.1169-1174.
- [14] V. Emel'yanenko, S. Verevkin. Thermodynamic properties of caffeine: Reconciliation of available experimental data. *J. Chem. Thermodynamics*. **2008**. No.40. P.1661-1665.
- [15] Jia-Xin Dong, Qiang Li, Zhi-Cheng Tan, Zhi-Heng Zhang, Yi Liu. The standard molar enthalpy of formation, molar heat capacities, and thermal stability of anhydrous caffeine. *J. Chem. Thermodynamics*. **2007**. No.39. P.108-114.
- [16] X.U. Fen, Q.IU. Shu-Jun, LIANG Jian-Guo, WU Rui-Hua, SUN Li-Xian, L.I. Fen. Low temperature heat capacity and thermal analysis of caffeine, theophylline and aminophylline. *Acta Phys. Chim. Sin.* **2010**. No.26(8). P.2096-2102.
- [17] F.N. Shamsetdinov, S.A. Bulaev, Z.I. Zaripov. Thermophysical properties of supercritical carbon dioxide – caffeine system. *Heruld of Kazan State Technological University*. **2010**. Vol.11. P.465-468. (russian)
- [18] U.K. Deiters, S.L. Randzio. A combined determination of phase diagrams of asymmetric binary mixtures by equations of state and transitiometry. *Fluid Phase Equilib.* **2007**. No.260. P.87-97.
- [19] J.H. Stern, E. Lowe. Enthalpies of transfer of theophylline and caffeine from water to aqueous alcohols at 25 °C. *Journal of Chemical and Engineering Data*. **1978**. Vol.23. No.4. P.341-342.
- [20] J.H. Stern, L.R. Beeninga. Partial molal heat capacities of caffeine and theophylline in pure water. *The Journal of Physical Chemistry*. **1975**. Vol.79. No.6. P.582-584. (russian)
- [21] Ingrid Stokkeland, Arne Skauge, Harald Hotiland. Changes in Partial Molar Volume and Isentropic Partial Molar Compressibility of Self-Association of Purine and Caffeine in Aqueous Solution at 1-1600 Bar. *Journal of Solution Chemistry*. **1986**. Vol.16. No.1. P.45-53.
- [22] C.S. Su., Y.P. Chen. Correlation for the solubility of pharmaceutical compounds in supercritical carbon dioxide. *Fluid Phase Equilibrium*. **2007**. No.254. P.167-173.
- [23] M. Johannsen, G. Brunner. Solubility of the xanthines caffeine, theophylline and theobromine in supercritical carbon dioxide. *Fluid Phase Equilibrium*. **1994**. No.95. P.215-260.
- [24] Azevedo Álvaro Bandeira Antunes de et al. Supercritical CO₂ recovery of caffeine from green coffee oil: new experimental solubility data and modeling. *Quím. Nova*. **2008**. Vol.31. No.6. P.1319-1323.
- [25] F.N. Shamsetdinov, Z.I. Zaripov and al. High yield biofuel production from vegetable oils with supercritical alcohols. *Liquid fuels: types, properties and production*. Nova Science Publishers, NY. **2012**.
- [26] R.A. Usmanov, F.N. Shamsetdinov, R.R. Gabitov, Sh.A. Biktashev, F.M. Gumerov, F.R. Gabitov, Z.I. Zaripov, R.A. Gazizov, R.S. Yarullin, I.A. Yakushev. Pilot plant for the continuous transesterification of vegetable oils in supercritical methanol, ethanol. *Supercritical Fluids: Theory and Practice*. **2011**. Vol.2. P.1-19. (russian)
- [27] I.A. Vasilyev, V.M. Petrov. Thermodynamic properties of oxygen-containing organic compounds. *Chemistry*. Leningrad. **1984**. 240p. (russian)
- [28] Ya.M. Naziyev, A.N. Shahverdiev, M.M. Bashirov, N.S. Aliyev. Thermo-properties of single-atom esters (isobaric heat capacity). *TVT*. **1994**. Vol.32. No.6. P.936. (russian)

- Full Paper** _____ L.N. Maskaeva, V.F. Markov, I.V. Vaganova, and N.A. Forostyanaya
- [29] Z.I. Zaripov, S.A. Burtsev, A.V. Gavrilov, S.A. Bulaev, G.Kh. Mukhamedzyanov. Thermal and calorimetric properties of *n*-butyl ester. *Heruld of Kazan State Technological University*. **2002**. Vol.1-2. P.208-212. (russian)
- [30] L.Y. Yarullin, F.M. Gumerov, Truong Nam Hung, I.I. Gilmutdinov, Z.I. Zaripov, F.R. Gabitov, A.B. Remizov. The composition and structure of the tea leaves, processed in supercritical carbon dioxide. *Butlerov Communications*. **2016**. Vol.48. No.11. P.88-100. ROI: jbc-02/16-48-11-88
- [31] Z.I. Zaripov, L.Yu. Yarullin, F.M. Gumerov, Truong Nam Hung, F.R. Gabitov, V.A. Kayumova. Thermal effects dissolution main tea components in supercritical carbon dioxide. *Butlerov Communications*. **2017**. Vol.49. No.2. P.96-101. P.96-101. ROI: jbc-02/17-49-2-96