

Investigation of the coal-similar wood by the methods of IR- and EPR-spectroscopy

© Ludmila S. Kocheva,^{1,3} Anatoly P. Karmanov,^{2,3*+} Vladimir P. Lutoev,¹
Igor Kh. Shumilov,¹ and Yury V. Glukhov¹

¹ Chemistry Mineral Raw Material Laboratory. Mineralogy Laboratory. Mineral-Raw Resources Laboratory. Institute of Geology. Komi Scientific Center. Ural Division. Russian Academy of Sciences. Pervomajskaja St., 54. Syktyvkar, 167982. Komi Republic. Russia.

Phone: +7 (8212) 24-54-16. E-mail: lskocheva@geo.komisc.ru

² Biochemistry and Biotechnology Laboratory. Institute of Biology. Komi Scientific Center. Ural Division. Russian Academy of Sciences. Kommunisticheskaya St., 28. Syktyvkar, 167982. Komi Republic. Russia.

Phone: +7 (909) 120-81-63. E-mail: apk0948@ib.komisc.ru

³ Chemistry Department. Pitirim Sorokin Syktyvkar State University. Oktiabrskii Av., 55. Syktyvkar. 167001. Komi Republic. Russia.

*Supervising author; +Corresponding author

Keywords: coal-similar wood, IR-spectroscopy, EPR-spectroscopy, lignin, activated carbon.

Abstract

By the IR- and EPR-pectroscopy methods it is carried out the study of organic matter of the coal-similar wood, which relates to the Devonian (CSW-1) and the Jurassic (CSW-2) geological periods. Element composition of the sample CSW-1, %: C – 74.4, H – 4.2, O – 20.2, N – 1.2; composition of the sample CSW-2, %: C – 54.4, H – 4.3, O – 39.7, N – 0.5. In the IR-spectra of the coal-similar wood samples the absorption bands with the maximums with 820-830, 860-870, ~1285, 1350-1370, 1445-1450, ~1610, 1650-1660, 2920-2940, 3380-3440 cm^{-1} present, which testify about the structural units of aromatic nature. The comparative analysis of the spectra of investigated samples, activated carbon, and also the moss *Polytrichum* lignin, the birch wood lignin and the fir tree wood lignin made it possible to make a conclusion about the similarity of organic matter of the coal-similar wood and contemporary lignins. It is established that the investigated samples possess the clearly expressed paramagnetic properties. EPR-signals are isotropic singlets with the g-factor, which corresponds to phenoxyl radicals (2.003-2.004), being present in the native lignins. The parameters of carbonic radicals correspond to the degree of the metamorphic transformations (carbonate formation) of initial organic matter, which is low for the sample CSW-2 and high of the sample CSW-1. The concentration of the free radicals C_{pmc} in the Devonian coal-similar wood samples is close to the limiting value for the carbonaceous substances. The presence of mineral phases is revealed by the EPR-spectroscopy method in the investigated samples. In the spectrum of CSW-2 sample the line in the region $g = 4.3$, caused by the high-spin state of the Fe^{3+} ions, is fixed. For the sample CSW-1 it are established the lines with $g = 2.29$ and 2.89 , which can be related to ions Fe^{3+} in the low-spin state. Intensive partial spectrum with the complex hyperfine structure, characteristic for vanadyl ions VO^{2+} is revealed for the sample CSW-2. As a result conducted investigations it is made conclusion about the proximity of organic matter of the coal-similar wood of Devonian and Jurassic geological periods to lignin's type polymers.

References

- [1] V.I. Peresyphkin, E.A. Romankevich. Biogeochemistry of lignin. Moscow: GEOS. 2010. 340p. (russian)
- [2] S. Dutta, S. Bhattacharya, M. Mallick, A.C. Shukla, U. Mann. Preserved lignin structures in Early Eocene Surat lignites, Cambay Basin, Western India. *Journal Geological Society of India*. 2012. Vol.79. P.345-352.
- [3] I.H. Shumilov, O.P. Telnova. Stage of lithogenesis Devonian deposits of middle Timan (of TSIL'ma river basin). *Bulletin of the Institute of Geology, Komi science center URD RAS*. 2014. No.4. P.16-19. (russian)
- [4] J.M. Pepper, P.E. Baylis, E. Adler. The isolation and properties of lignin obtained by the acidolysis of spruce and aspen woods in dioxane-water. *Canad. J. Chem.* 1959. Vol.37. No.8. P.1241-1245.
- [5] B.K. Saikia, R.K. Boruah, P.K. Gogoi. FT-IR and XRD analysis of coal from Makum coalfield of Assam. *J. Earth Syst. Sci.* 2007. Vol.116. No.6. P.575-579.

- Full Paper** _____ L.S. Kocheva, A.P. Karmanov, V.P. Lutoev, I.Kh. Shumilov, Yu.V. Glukhov
- [6] A.P. Karmanov, L.S. Kocheva, M.F. Borisenkov. Chemical composition and sorption properties of sorbents based on lignin. *Butlerov Communications*. **2016**. T.45. No.1. P.76-84. DOI: 10.37952/ROI-jbc-01/16-45-1-76
- [7] A.V. Canary, Z.A. Canary, L.S. Kocheva, A.P. Karmanov, N.I. Karlin, O.A. Konyk. The Chemical structure of lignins and their sorption capacity in relation to the mycotoxin T-2. *Butlerov Communications*. **2016**. Vol.46. No.5. P.67-73. DOI: 10.37952/ROI-jbc-01/16-46-5-67
- [8] F.A. Ants, V.M. Vinokurov, A.A. Galeev, G.R. Bulka, N.M. Nizamutdinov, N.M. Khasanova. Paramagnetism and nature of dispersed organic matter in the Permian deposits of Tatarstan. *Georesurs*. **2006**. Vol.2. No.19. P.40-45 (russian)
- [9] J.E. Conard. In: Magnetic Resonance. Introduction, Advanced Topics and Application to Fossil Energy. *Dordrecht: Reidel*. **1984**. P.441-459.