

## The dumbbell model of the helium atom

© Alexey A. Potapov

Butlerov Science Foundation. Bondarenko, St. 33-44. Kazan, 420066. Russia.

Phone: +7 (843) 231-42-30; +7 (395) 246-30-09. E-mail: [aleksey.potapov.icc@gmail.com](mailto:aleksey.potapov.icc@gmail.com)

**Keywords:** helium atom, elliptical orbits, shell model, structure.

### Abstract

The purpose of this article is to discuss the problems of the electronic structure of a helium atom. Given the evolution of ideas about the structure of the helium atom. The first physically reasonable model of the helium atom was the atom shell model, proposed N. Bohr. Changing this to closure shell model came quantum-mechanical  $s^2$ -spherical model of the electron cloud. In the quest to improve shell Bohr's model the author proposes a model with two elliptical orbits with common focus on the nucleus of the atom. This article proposes for discussion a revised model of the dumbbell structure of the helium atom. In this model, the charge of the cation  $\text{He}^+$  acts as the center of attraction of valence electrons and carries out the capture of an electron on elliptical orbit and the education two-elliptical shell. The shell of the atom is a system of two coupled elliptic orbits formed mirror symmetrical relative to the core electrons. In this configuration, the elliptical orbits of the atom are relatively independent. A distinctive feature of this two-elliptical of the model is that when calculating the eccentricity of elliptical orbits was used the energy of the circular orbit, which was determined based on finding the minimum of the potential function,  $\epsilon_m = 27.5$  eV. The main parameters of a helium atom: the semi-major axis  $a \approx 0.415$  Å, while the semi-minor axis at  $b \approx 0.37$  Å, eccentricity  $\vartheta = 0.32$ , the minimum  $r_1 = 0.28$  Å and maximum  $r_2 = 0.55$  Å, corresponding to the distance of the apogee  $h_a$  and the effective radius  $a_{\text{eff}}$ , so  $r_2 = r_A = a_{\text{eff}}$ . For comparison, according to the literature, the radius of a helium atom is in the range  $a \approx 1.2$ - $1.4$  Å. An indirect confirmation of the proposed model can serve as: the fact of the crystallization of helium, the doublet nature of the optical spectrum of a helium atom. In the framework of the dipole shell model dumbbell model of a helium atom became the basis for the understanding of the mechanism of formation of multi-electron atoms. The helium atom can serve as next example of rightness the classical approach in atomic physics.

### References

- [1] N. Bor. Selected Scientific Works. *Moscow: Science*. **1970**. 584p. (russian)
- [2] P. Atkins. Physical chemistry. Vol.1. *Moscow: Mir*. **1980**. 584p. (russian)
- [3] E.V. Shpolsky. Atomic physics. *Moscow: Fizmatgiz*. **1963**. Vol.1. 576p. (russian)
- [4] Physical Encyclopedic Dictionary. *Moscow: Soviet encyclopedia*. **1983**. 928p. (russian)
- [5] R. Dickerson, G. Gray, J. Haight. Basic laws of chemistry. Vol.1. *Moscow: Mir*. **1982**. 652p. (russian)
- [6] U. Fleiger. Structure and dynamics of molecules. Vol.1. *Moscow: Mir*. **1982**. 408p. (russian)
- [7] A.A. Potapov. Renaissance of the classical atom. *Moscow: Publishing House "Science"*. LAPLAMBERTA academic publishing. **2011**. 444p. (russian)
- [8] A.A. Potapov. Fundamental bases of the structure of substance. *Butlerov Communications*. **2015**. Vol.41. No.2. P.1-29. ROI: jbc-02/15-41-2-1
- [9] A.A. Potapov. Science about substance: a condition of researches. *Butlerov Communications*. **2011**. Vol.24. No.1. P.1-45. ROI: jbc-02/11-24-1-1
- [10] Potapov A.A. Science of matter: a way out of crisis. On Sat "Actual problems of biology, chemistry, physics": materials of the international scientific-practical conference. *Novosibirsk: Publishing house "EKOR-book"*. **2011**. P.136-148. (russian)
- [11] A.A. Potapov. High technologies: substance control algorithm. *LAPLAMBERTA academic publishing*. **2016**. 122p. (russian)
- [12] I.E. Tamm. Fundamentals of the theory of electricity. *Moscow: Science*. **1976**. 616p. (russian)
- [13] Ch. Kittel, V. Knight, M. Ruderman. Mechanics. *Moscow: Science*. **1983**. 448p. (russian)
- [14] S.S. Batsanov. Structural chemistry. Facts and dependencies. *Moscow: Dialogue-MSU*. **2000**. 292p. (russian)
- [15] M.I. Shahparonov. Introduction to the modern theory of solutions. *Moscow: High School*. **1976**. 296p. (russian)
- [16] Chemical encyclopedia: in 5 tons: v.1. *Moscow: Sov.entsikl*. **1988**. 623p. (russian)

- [17] A.A. Potapov. Noble gases: structure and properties: monogr. *Science today: theory, practice, innovation*. **2015**. Vol.7. P.72-177. (russian)
- [18] A.A. Potapov. The nature and mechanisms of binding atoms. *Moscow: RIOR: INFRA-M*. **2013**. 295p. (russian)
- [19] A.A. Potapov, Yu.V. Mineev. Planetary model of the hydrogen atom and hydrogen-like structures. *Butlerov Communications*. **2015**. Vol.44. No.11. P.1-15. ROI: jbc-02/15-44-11-1
- [20] A.A. Potapov. The radius of the hydrogen atom: a fundamental constant. *Science, technology and education*. **2015**. No.10. P.7-16. (russian)