

## Biomass production of carotenoid synthesizing yeasts of genus *Rhodotorula* at cultivation on agricultural wastes

© Olga P. Chervyakova<sup>+</sup>, Nikolay A. Suyasov\*, Alexandra M. Fomicheva, Irina V. Shakir, and Viktor I. Panfilov

Department of Biotechnology. D. Mendeleev University of Chemical Technology of Russia. Miusskaya sq. 9. Moscow, 125047, Russia. Phone: +7 (965) 362-99-17. E-mail: chervyakova85@mail.ru

\*Supervising author; <sup>+</sup>Corresponding author

**Keywords:** carotenoids, *Rhodotorula*, agricultural wastes, plant raw materials, feed supplement.

### Abstract

Carotenoids are biologically active substances that play an important role in the course of physiological processes in humans and animals. Besides to plants, many microorganisms are able to synthesize carotenoid pigments, therefore they are used as producers in the industrial production of carotenoids. The yeasts of genus *Rhodotorula* can synthesize carotenoids, the cultivation of which can be carried out using various sources of carbon, including components of plant raw materials. The possibility of using secondary plant raw materials (by-products of the agricultural and food industry, substandard raw materials) allows solving the problems of their processing, that is actual at the present time.

This research work is devoted to evaluating the bio-potential of various types of plant raw materials for the cultivation of carotenoid synthesizing yeasts *Rhodotorula rubra*.

It is found that pretreatment of plant raw materials, as well acid hydrolysis as enzymatic, allows increasing the availability of nutrient components, while the content of total carbohydrates in hydrolysates reaches 30.0-40.0 g/L. The degree of assimilation of carbohydrates by yeast *Rhodotorula rubra* is 60.0-80.0%, the accumulation of biomass varies in the range from 8.0 to 10.5 g/L.

Analysis of the carotenoid composition of the biomass showed that the amount of pigments and their ratio varies depending on the type of substrate, on the method of its processing, and ranges from 50.0 to 550.0 µg/L. The greatest potential for microbial synthesis of carotenoids by yeast *Rhodotorula rubra* is beet molasses and acid hydrolysates of deproteinised sunflower meal and of Jerusalem artichoke tuber, the maximum yield of pigments from a liter of culture fluid being 507.0, 545.0 and 180.3 µg, respectively.

### References

- [1] G.I. Frengova, D.M. Beshkova. Carotenoids from *Rhodotorula* and *Phaffia*: yeasts of biotechnological importance. *Journal of Industrial Microbiology & Biotechnology*. **2009**. Vol.36. No.2. P.163.
- [2] E. Valduga, A. Valério, H. Treichel, M. Di Luccio, A. Furigo Júnior. Study of the bioproduction of carotenoids by *Sporidiobolus salmonicolor* (CBS 2636) using pretreated agroindustrial substrates. *Journal of Chemical Technology and Biotechnology*. **2008**. Vol.83. No.9. P.1267-1274.
- [3] M.M.K. Bagy, M.H. Abd-Alla, N.A. Nafady, F.M. Morsy, G.A.E. Mahmoud. Bioconversion of plant wastes to β-carotene by *Rhodotorula glutinis* KU550702. *European Journal of Biological Research*. **2016**. Vol.6. No.4. P.226-241.
- [4] S. Petrik, S. Obruča, P. Benešová, & I. Márová. (2014). Bioconversion of spent coffee grounds into carotenoids and other valuable metabolites by selected red yeast strains. *Biochemical Engineering Journal*. **2014**. Vol.90. P.307-315.
- [5] T. Schneider, S. Graeff-Hönninger, W.T. French, R. Hernandez, N. Merkt, W. Claupein. Lipid and carotenoid production by oleaginous red yeast *Rhodotorula glutinis* cultivated on brewery effluents. *Energy*. **2013**. Vol.61. P.34-43.
- [6] O.P. Chervyakova, I.V. Shakir, N.A. Suyasov, V.I. Panfilov. Factors affecting carotenoid biosynthesis by yeast *Rhodotorula rubra*. *Chemical Industry Today*. **2015**. No.5. P.45-50. (russian)
- [7] E.V. Bashashkina, I.V. Shakir, N.A. Suyasov, V.I. Panfilov. Use of coffee sludge as a raw material for production of protein feed supplement. *Chemical Industry Today*. **2010**. No.6. P.28-33. (russian)
- [8] E.V. Bashashkina, N.A. Souyasov, I.V. Shakir, V.I. Panfilov. Bioconversion of soluble coffee production waste into fodder products. *Ecology & Industry of Russia*. **2011**. No.1. P.18-19. (russian)

- Full Paper** \_\_\_\_\_ O.P. Chervyakova, N.A. Suyasov, A.M. Fomicheva, I.V. Shakir, and V.I. Panfilov
- [9] B.A. Karetkin, T.S. Kataeva, D.V. Baurin, V.D. Grosheva. Factorial design for optimization of media pretreatment conditions. *Fundamental Research*. **2014**. No.11-1. P.13-19. (russian)
- [10] Dr. Karetkin B.A., Chervyakova O.P., Dr. Baurin D.V., Assoc. Prof. Dr. Shakir I.V., Prof. Dr. Panfilov V.I. Optimisation of pretreatment condition of plant raw composed media for carotenoids biosynthesis by yeast *Rhodotorula rubra*. *16th International Multidisciplinary Scientific GeoConference, SGEM 2016*. **2016**. P.301-309.
- [11] A.S. Vecher, A.N. Kulikova. Spectrophotometric determination of the content of carotenoids in the biosynthesis of microorganisms. *Physiological and biochemical studies of plants. Minsk*. **1967**. P.46-54.