

***N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine as inhibitor of radical chain oxidation of ethylbenzene**

© **Gulnaz M. Sharipova,⁺¹ Irina V. Safarova,¹ Anatoly Ya. Gerchikov,^{1*+} Ildus Sh. Nasyrov,² and Gulfina N. Musina¹**

¹*Department of Physical Chemistry and Chemical Ecology. Bashkir State University.*

Z. Validi St., 32. Ufa, 450076. Republic of Bashkortostan. Russia.

Phone: +7 (347) 273-67-27. Fax: +7 (347) 273-67-01.

E-mail: gerchikov@inbox.ru, gulnaz-sharipova@list.ru⁺

²*AO «Syntez Cauchuk». Tekhnicheskaya St., 14. Sterlitamak, 453110. Russia.*

*Supervising author; ⁺Corresponding author

Keywords: *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine, antioxidants, kinetics, activation parameters.

Abstract

The problem of stabilization of organic materials has important practical value. For the purpose of protection of synthetic rubbers against oxidation decomposition in the production enterprises inhibitors of aminny and phenolic types are used. For selection of efficient stabilizers – antioxidants it is necessary to know the quantitative characteristics of their effectiveness. One of important quantitative characteristics of antioxidants is the kinetic constant of break of chain on an inhibitor at oxidation of an organic substratum. On the example of model reaction of the initiated oxidation of ethylbenzene anti-oxidizing properties of *N*-2-ethylhexyl-phenyl-*p*-phenylenediamine are studied. The kinetics of reaction of the initiated oxidation of ethyl benzene in the presence of additives *N*-2-ethylhexyl-phenyl-*p*-phenylenediamine was studied with use of highly sensitive manometric installation. It is shown that introduction to the model system of the studied substance leads to decrease of saturating speed of oxygen. Besides, on the kinetic curves of oxygen uptake received at measurement of saturating speed of oxygen in the presence of various concentration *N*-2-ethylhexyl-phenyl-*p*-phenylenediamine induction periods are observed, that is characteristic of the strong inhibitors, to which number it is possible to carry also studied inhibitor. As a result a research the quantitative kinetic parameters characterizing anti-radical activity of substance in the form of a kinetic constant of break of chain of oxidation and capacity of an inhibitor are received. Activation parameters of interaction of connection with peroksilny radicals of ethyl benzene are determined that allows to predict effectiveness of anti-oxidizing effect of the studied antioxidant depending on temperature.

References

- [1] Yu.V. Vinokurov, S.M. Kavun, N.I. Koltsov. Development of process of receiving *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamines on the basis of a research of features of his synthesis in laboratory and industrial conditions. *Bulletin of the Kazan Technological University*. **2011**. No.15. P.48-56. (russian)
- [2] *Patent. 2293077 Russian Federation, MPK C07C211/51, C07C 211/55.* Way of receiving *N*-alkyl-*N*1-phenyl or *N*, *N*1-dialkil-of *p*-phenylenediamines. M.S. Yaskova, Yu.V. Vinokurov, V.B. Vyazova, A.I. Efremov, V.M. Filippov, S.M. Kavun. № 2005123571/04; request. 25.07.2005; publish. 10.02.2007. (russian)
- [3] *Patent 2362767 Russian Federation, MPK C07C211/51, C07C211/55, C07C209/16.* Way of receiving *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine. Yu.V. Vinokurov, V.M. Filippov, S.M. Kavun, V.A. Ivanov, M.S. Yaskova. № 2007147309/04; request. 18.12.2007; publish. 27.07.2009. (russian)
- [4] Yu.V. Vinokurov Perfecting of the production technology of the stabilizer of rubbers and rubbers *n*-2-ethylhexyl-*n'*-phenyl-*p*-phenylenediamine: The thesis for a degree of the candidate of technical sciences. *Kazan*. **2012**. (russian)
- [5] E.T. Denisov, N.I. Mickiewicz, V.E. Agabekov. Mechanism of liquid-phase oxidation of oxychemicals. *Minsk: Science and Technology*. **1975**. 335p.
- [6] N.M. Emanuel, D. Gaul. Ethyl benzene oxidation. *Moscow: Nauka*. 1984. 376p. (russian)

- [7] N.M. Emanuel, E.T. Denisov, Z.K. Mayzus. Chain reactions of oxidation of hydrocarbons in a liquid phase. *Moscow: Nauka*. **1965**. (russian)
- [8] A. Weisberg, E. Proskauer, J. Riddik, E. Tups. Organic solvents. Physical properties and methods of cleaning. *Moscow: Publishing House of Foreign Literature*. **1958**. P.294. (russian)
- [9] E.T. Denisov. Kinetic constants of homolytical liquid-phase reactions. *Moscow: Nauka*. **1971**. (russian)
- [10] E.T. Denisov, V.V. Azatyan. Ingibirovaniye of chain reactions. *Chernogolovka: IFHCH OF RAS*. **1997**. 268p. (russian)