

Amid derivatives of salicylic acid – effective inhibitors of UV initiated oxidation of organic substrates

© Nadezhda M. Storozhok,^{*†} and Nadezhda P. Medyanik

Tyumen State Medical University Ministry of Health Russia. Odesskaya St., 54. Tyumen, 625023. Russia.

Fax: +7 (3452) 20-74-21. E-mail: nadinstor@mail.ru

^{*}Supervising author; [†]Corresponding author

Keywords: salicylic acid amide, antioxidant, anti-radical activity, UV-radiation.

Abstract

The results of the research are presented kinetics of UV initiated oxidation model substrate (methyloleat) in the presence of individual amides: 2-hydroxybenzoic acid *N*-(4-hydroxyphenyl)amide (**I**); 2-hydroxy-3-tretbutyl-5-ethylbenzoic acid *N*-(4-hydroxy-3,5-di-tret-butylphenyl)amide (**II**); 2-hydroxybenzoic acid *N*-[3-(4-hydroxy-3,5-di-tret-butylphenyl)prop-1-yl]amide (**III**); 2-hydroxy-3-tretbutyl-5-ethylbenzoic acid *N*-[3-(4-hydroxy-3,5-di-tret-butylphenyl)prop-1-yl]amide (**IV**) in comparison with reference antioxidants – dibunol and α -tocopherol. It is shown that all amides of salicylic acid **I-IV** effectively inhibit the process UV initiated oxidation of methyloleat. It is studied a relationship between antioxidant properties of amides salicylic acid **I-IV** of UV initiated oxidation and features of their structure. It is established that introduction shielding ortho-*t*-butyl substituents and separation of aromatic fragments by three methylene groups leads to a significant increase of antioxidant activity. It is shown that amide **IV** in the concentration range $(0.5-2.0) \cdot 10^{-4}$ Mol/L exceeds the efficiency of dibunol by 20%.

Testing the anti-radical activity of amides **I-IV** estimated by the chemiluminescence method from the constant rate of the reaction with peroxy radicals allowed to establish the range of changes in value $k_7 = 0.52 - 6.86 \cdot 10^4$, mol⁻¹·s⁻¹. It is shown that screening of phenolic OH-groups tret-butyl substituents and reduction of conjugation of electron density in molecules by separating the residue of salicylic acid and phenol by methylene groups leads to a significant decrease in anti-radical activity of antioxidants. Thus, the largest value of k_7 was established for the amide **I**, which in its structure has two unsubstituted phenolic hydroxyls. The antiradical activity of the shielded analogue (amide **II**) is 4 times inferior. The mechanism of the inhibitory action of amides of salicylic acid was established, it includes: direct interaction of phenols with free radicals; destruction of hydroperoxides of lipids due to the amide fragment of the molecule; partial absorption of UV radiation, associated with the presence of a salicylic acid residue.

References

- [1] N.M. Emanuel, E.T. Denisov, Z.K. Mayzus. Chain reactions of oxidation of hydrocarbons in the liquid phase. *Moscow: Science*. **1966**. 375p. (russian)
- [2] V.Ya. Shlyapintok. Photochemical transformations and stabilization of polymers. *Moscow: Chemistry*. **1979**. 344p. (russian)
- [3] B.N. Gorbunov, Ya.A. Gurvich, I.P. Maslova. Chemistry and technology of stabilizers of polymeric materials. *Moscow: Chemistry*. **1981**. 368p. (russian)
- [4] M. Diepens, P. Gijsman. Photodegradation of bisphenol a polycarbonate with different types of stabilizers. *Polymer Degradation and Stability*. **2010**. Vol.95. No.5. P.811-817.
- [5] T.G. Denisova, E.T. Denisov. Reactivity of natural phenols in radical reactions. *Kinetics and Catalysis*. **2009**. Vol.50. No.3. P.335-343.
- [6] Oxidative stress. Prooxidants and antioxidants. E. B. Menschikova [et al]. *Moscow: Firm "Word"*. **2006**. 554p. (russian)
- [7] Phenolic bioantioxidants. N. K. Zenkov [et al]. *Novosibirsk*. **2003**. 328p. (russian)
- [8] Features of inhibitory action of antioxidants of the IBCPhN group. N.M. Storozhok, M.G. Perevozkina. *Scientific Bulletin of the Tyumen Medical Academy*. **2003**. No.1. P.52-59. (russian)
- [9] To the mechanism of action of new sulfur-containing antioxidants. N.M. Storozhok, N.V. Gureeva, A.P. Krysin. *Scientific Bulletin of the Tyumen Medical Academy*. **2003**. No.1. P.18-22. (russian)

- AMID DERIVATIVES OF SALICYLIC ACID – EFFECTIVE INHIBITORS OF UV INITIATED OXIDATION... __ 22-29*
- [10] Synthesis, structure and antioxidant activity of sulfur tetrakispenol G.N. Nugumanova [et al]. *Journal of Organic Chemistry*. **2010**. Vol.80. No.7. P.1175-1184. (russian)
- [11] Synthesis, structure and activity of antioxidants "hybrid structure" based on N-substituted amides of salicylic acid. N.M. Storozhok, N.P. Medyanik, A.P. Krysin et al. *Journal of Organic Chemistry*. **2013**. Vol.49. No.7. P.1046-1049. (russian)
- [12] Photolysis of sulfosalicylic acid in aqueous solutions over a wide pH range. I. P. Pozdnyakov [et al]. *J. Photochem. Photobiol. A: Chem.* **2006**. Vol.181. No.1. P.37-43.
- [13] Determination of rate constants and inhibition coefficients of phenol antioxidants using a model chain reaction. V.F. Tsepalov [et al]. *Kinetics and Catalysis*. **1977**. Vol.18. No.5. P.1261-1267. (russian)
- [14] Chemiluminescent methods for studying slow chemical processes. V.Ya. Shlyapintokh [et al]. *Moscow: Science*. **1966**. 138p. (russian)
- [15] About the role of the elementary reaction 10 in the kinetics of the oxidation of polyunsaturated lipids. N.M. Storozhok, N.V. Gureeva. *Scientific Bulletin of the Tyumen Medical Academy*. **2003**. No.1. P.27-35. (russian)
- [16] N.M. Storozhok, N.P. Medyanik, S.A. Krekov, and A.P. Krisin. The supramolecular organization N-replaced amides of salicylic acid. *Butlerov Communications*. **2011**. Vol.24. No.3. P.109-112. ROI: jbc: 02/11-24-3-109
- [17] V.L. Antonovsky. Organic peroxy initiators. *M.: Chemistry*. **1972**. 448p. (russian)
- [18] Kinetics and mechanism of photochemical reactions of N-substituted amide of salicylic acid. N.P. Medyanik, N.V. Gureeva, N.M. Storozhok. *News of the Tomsk Polytechnic University*. Vol.318. No.3. P.116-120. (russian)
- [19] Photochemistry of N-substituted salicylic acid amides. I.P. Pozdnyakov, N.M. Storozhok, N.P. Medyanik et al. *Russian Chemical Bulletin, International Edition*. **2015**. Vol.64. No.6. P.1319-1326.