

The reaction of *N*-aryl-2,5-dihydropyrrol-2,5-diones with α -furylmethanol

© Yury N. Mitrasov,^{1*} Anzhelika A. Avruyskaya,¹
Oksana V. Kondratieva,¹ and Vadim A. Kozlov³

¹ Department of Bioecology and Chemistry. Chuvash State Pedagogical University. I. Yakovlev. K. Marks St., 38. Cheboksary, 428000. Chuvash Republic. Russia.
Phone: +7 (8352) 45-68-45. E-mail: mitrasov_un@mail.ru.

² Department of Medical Biology with Course of Microbiology and Virology. Chuvash State University. I. N. Ulyanov. Moskovskiy pr., 15. Cheboksary, 428015. Chuvash Republic. Russia.
Phone: +7 (8352) 45-26-97. E-mail: pooh12@yandex.ru

*Supervising author; ⁺Corresponding author

Keywords: α -furylmethanol, *N*-aryl-2,5-dihydropyrrol-2,5-diones, diene synthesis, fluorescence.

Abstract

The article describes the interaction of the isomeric *N*-tolyl-*N*-1-naphtyl-2,5-dihydropyrrol-2,5-diones with α -furylmethanol. The research urgency is caused by the fact that 2,5-dihydropyrrol-2,5-dione (imide maleic acid) and their derivatives represent a promising class of organic compounds. Due to the presence of highly reactive double bonds they can easily polymerized and are copolymerized with various unsaturated monomers, interact with a variety of nucleophilic reagents (primary and secondary amines, cyanate, isocyanates, azomethine, epoxides, etc.), participate in a number of cycloaddition reactions. For example, the Diels-Alder in the presence of suitable dienes, reaction with double bonds of allyl type. A number of derivatives of maleinimide finds application as pesticides. Thus, they exhibit a high insecticidal, fungicidal and herbicidal activity. In addition, maleinimide proposed as highly effective pharmaceuticals for the treatment of several diseases, including cardiovascular, Alzheimer's, diabetes of the 2nd type, cancer and HIV. Reaction of isomeric *N*-tolyl-, *N*-benzyl- and *N*-1-naphtyl-2,5-dihydropyrrol-2,5-diones with α -furylmethanol carried out at equimolar ratio of reagents at room temperature in absolute 1,4-dioxane or benzene. Methods of gas chromatography-mass spectrometry, IR and ¹H NMR spectroscopy established that the reactions proceed according to the scheme of [4+2]-cycloaddition with the formation of 4-*N*-aryl-1-hydroxymethyl-3,5-dioxo-10-oxa-4-azatricyclo[5.2.1^{1.7}.0^{2.6}]dec-8-enes. The possibility of education in the molecules synthesized adducts intramolecular hydrogen bonds involving carbonyl and hydroxyl groups. Adducts of diene synthesis are promising objects for the study of fluorescent activity. To study the phenomenon called fluorescence, the crystals of the samples were subjected to microscopy. It was revealed that they observed three peaks of fluorescence at 507, 534 and 569 nm, which corresponds to yellow-green region of the spectrum.

References

- [1] O.A. Kolyamshin, V.A. Danilov, S.Yu. Vasilieva. Maleinimide: synthesis, properties and biological activity. *Dep. VINITI*, No. 250-B2010 from 06.05.2010. 75p. (russian)
- [2] Yu.N. Mitrasov, A.A. Avruyskaya, L.M. Sadikova, O.V. Kondratyeva, O.A. Kolyamshin. The reaction of *N*-aryl-2,5-dihydropyrrol-2,5-diones with functional derivatives of furan. *Cheboksary: Chuvash. State Pedagogical University*. 2013. 155p. (russian)
- [3] O.A. Kolyamshin, Yu.N. Mitrasov, V.A. Danilov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted nitrogen-containing heterocycles. Part 1. Synthesis of *gem*-dichlorocyclopropylmethyltiosubstituted derivative benzimidazoles, benzoxazoles, 4,6-dimethylpyrimidines and 1,2,4-triazoles. *Butlerov Communications*. 2016. Vol.45. No.3. P.6-10. DOI: 10.37952/ROI-jbc-01/16-45-3-6
- [4] O.A. Kolyamshin, Yu.N. Mitrasov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted nitrogen-containing heterocycles. Part 2. Synthesis and properties of 1-(2,2-dichlorocyclopropylmethyl)pyrazoles and 5-pyrazolones. *Butlerov Communications*. 2016. Vol.46. No.4. P.77-80. DOI: 10.37952/ROI-jbc-01/16-46-4-77
- [5] O.A. Kolyamshin, Yu.N. Mitrasov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted

- Full Paper** _____ Yu.N. Mitrasov, A.A. Avruyskaya, O.V. Kondrat'eva, and V.A. Kozlov nitrogen-containing heterocycles. Part 3. Synthesis and properties of 3-[(1-methyl-2,2-dichlorocyclopropyl)methyl]hydantoins. *Butlerov Communications*. **2016**. Vol.48. No.10. P.103-107. DOI: 10.37952/ROI-jbc-01/16-48-10-103
- [6] V.A. Kozlov, Yu.N. Mitrasov, O.V. Kondrat'eva, I.V., Gordeeva, O.B. Polyakova, O.E. Kirillova, A.N. Kochneva, A.A. Avruyskaya, V.A. Sorokina, O.A. Kolyamshin. Photometry 4-Aza-4-aryl-1-hidroksimethyl-10-oxa-3,5-dioxotricyclo[5.3.1^{1.7}.0^{2.6}]dec-8-enes. *Materials of the seventh international scientific school "Science and innovations – 2012" ISS "SI-2012"*. Yoshkar-Ola: Mari State University. **2012**. P.232-235. (russian)
- [7] Yu.N. Mitrasov, A.A. Avruyskaya, O.V. Kondratieva. The interaction of the nitro- and aminosubstituted of phenylmaleimide and furfurylated alcohol. *Journal of General Chemistry*. **2015**. Vol.85. No.1. P.82-85. (russian)
- [8] Yu.N. Mitrasov, A.A. Avruyskaya, O.B. Polyakova, O.E. Ivanova. Interaction derivative (2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)benzoic acids α -furylmethanol. *Journal of Organic Chemistry*. **2015**. Vol.51. No.8. P.1206-1207. (russian)
- [9] Yu.N. Mitrasov, I.V. Gordeyeva, O.V. Kondratyeva, A.A. Avruyskaya, O.E. Kirillova. Reactions of derivatives of 2,5-dihydro-2,5-dioxo-1H-pyrrol-1-eleanorrigby (-phosphonic) acids and furfurylated alcohol. *Bulletin of the Chuvash State Pedagogical University of I.Ya. Yakovlev. Ser. natural and technical Sciences*. **2012**. No.2 (74). Part1. P.34-37. (russian)
- [10] V.A. Kozlov, Yu.N. Mitrasov, A.A. Avruyskaya, O.B. Polyakova, O.V. Kondratieva. Acute toxicity of derivatives of *m*- and *p*-(4-Aza-1-hidroksimethyl-10-oxa-3,5-dioxotricyclo[5.2.1^{1.7}.0^{2.6}]dec-8-en-4-yl)benzoic acids. *The ninth international scientific school "Science and innovations – 2014" ISS "SI-2014"*. Yoshkar-Ola: Mari State University. **2014**. P.183-185. (russian)