

Phenolic profile of *Thymus reverdattoanus* Serg., an endemic species of Sakha (Yakutia) flora: Chemotaxonomic aspect

© Nadezhda K. Chirikova,^{1*} Nina I. Kashchenko,² and Daniil N. Olennikov²⁺

¹ Laboratory of Biochemistry and Biotechnology. Institute of Natural Science. Ammosov's North-Eastern Federal University. Kulakovskogo St., 48. Yakutsk, 677007. Republic Sakha (Yakutia). Russia.

Phone: +7 (4112) 35-32-13. E-mail: hofnung@mail.ru

² Laboratory of Medical and Biological Research. Institute of General and Experimental Biology, Siberian Division, Russian Academy of Science. Sakh'yanyovoy St., 6. Ulan-Ude, 670047. Republic Buryatia. Russia.

Phone: +7 (3012) 43-45-43. E-mail: olennikovdn@mail.ru

*Supervising author; +Corresponding author

Keywords: *Thymus reverdattoanus*, Lamiaceae, endemic species, flavonoids, caffeic acid derivatives, HPLC, chemotaxonomy.

Abstract

Reverdatto's thyme (*Thymus reverdattoanus* Serg.) is an endemic species of the family Lamiaceae, common in the Arctic floristic region of the Republic of Sakha (Yakutia). For the first time we realized a study of the chemical composition of *T. reverdattoanus* herb using a complex of chromatographic and spectral methods which led to the discovery of 30 phenolic compounds. Among the phenylpropanoids, caffeic acid and its derivatives were identified as esters with quinic acid (3-*O*-caffeoylquinic acid and 1,3-di-*O*-caffeoylquinic acid), as well as dimeric and trimeric forms (lithospermic acid, rosmarinic acid, salvianolic acid B). The flavonoids identified (aglycons and glycosides) belonged to the groups of dihydroflavonols (taxifolin and its 3-*O*-glucoside), flavones (luteolin and its 6-*C*-glucoside, 7-*O*-rutinoside and 7-*O*-glucuronide, apigenin and its 7-*O*-rutinoside and 7-*O*-glucuronide) and flavanones (eriodictyol-7-*O*-rutinoside; naringenin and its 7-*O*-rutinoside; 7-*O*-neohesperidoside and 7-*O*-glucoside). The presence of isosakuranetin (4'-methoxy-naringenin) and its 7-*O*-glycosides, isosakuranine and didymine, was firstly shown for *T. reverdattoanus* and the genus *Thymus*. The lipophilic flavonoids including cirsimaritin, thymonin-4'-methyl ester and xanthomicrol were identified. Also, the presence of 3,4-dihydroxyphenyllactic acid, carvacrol and thymol was detected. Using the HPLC method, the quantitative content of the identified compounds was determined. It was shown that rosmarinic acid and luteolin-7-*O*-glucuronide were the main components of *T. reverdattoanus* with concentration levels 12.93 and 16.38 mg/g, respectively. The data obtained allowed to determine the chemotaxonomic significance of individual compounds for the species *T. reverdattoanus* and the genus *Thymus* at all. Some lipophilic flavonoids as well as widely represented in this plant glycosides of 5,7-dihydroxyflavanones have been identified as marker compounds of the species.

References

- [1] N.I. Kashchenko, N.K. Chirikova, D.N. Olennikov. Agrimoniin, an active ellagitannin from *Comarum palustre* herb with anti- α -glucosidase and antidiabetic potential in streptozotocin-induced diabetic rats. *Molecules*. **2017**. Vol.27. No.73.
- [2] D.N. Olennikov, N.K. Chirikova, Z.M. Okhlopkova, I.S. Zulfugarov. Chemical composition and antioxidant activity of *Tánara Otó* (*Dracocephalum palmatum* Stephan), a medicinal plant used by the North-Yakutian nomads. *Molecules*. **2013**. Vol.18. P.14105-14121.
- [3] D.N. Olennikov, N.K. Chirikova. Dracopalmaside, a new flavonoid from *Dracocephalum palmatum*. *Chem. Nat. Comp.* **2015**. Vol.51. P.1067-1069.
- [4] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova. Meadowsweet teas as new functional beverages: Comparative analysis of nutrients, phytochemicals and biological effects of four *Filipendula* species. *Molecules*. **2017**. Vol.27. No.16.
- [5] N.K. Chirikova, D.N. Olennikov. Chemodiversity and biological activity of sinantropic plants of Siberia. I. *Galeopsis bifida* Boenn. *Chemistry of Plant Raw Materials*. **2016**. No.2. P.230-237. (russian)

- [6] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova, L.M. Tankhaeva. Iridoids and flavonoids of four Siberian gentians: Chemical profile and gastric stimulatory effect. *Molecules*. **2015**. Vol.20. P.19172-19188.
- [7] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova. Bitter gentian teas: Nutritional and phytochemical profiles, polysaccharide characterisation and bioactivity. *Molecules*. **2015**. Vol.20. P.20014-20030
- [8] D.N. Olennikov, N.K. Chirikova. Algidisides I and II, new iridoids glycosides from *Gentiana algida*. *Chem. Nat. Comp.* **2016**. Vol.52. P.550-554.
- [9] D.N. Olennikov, N.K. Chirikova, C. Vennos. Chemical composition, antioxidant and anticholinesterase activities of *Gentianella azurea* from Russian Federation. *Nat. Prod. Commun.* **2017**. Vol.12. P.55-56.
- [10] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova. Spinacetin, a new caffeoylglucoside, and other phenolic compounds from *Gnaphalium uliginosum*. *Chem. Nat. Comp.* **2015**. Vol.51. P.1085-1090.
- [11] D.N. Olennikov, N.K. Chirikova. Caffeoylglucaric acids and other phenylpropanoids of the Siberian *Leonurus* species. *Chem. Nat. Comp.* **2016**. Vol.52. P.780-782.
- [12] D.N. Olennikov, N.K. Chirikova. Phlotuberosides I and II, new iridoids glycosides from *Phlomodites tuberosa*. *Chem. Nat. Comp.* **2017**. Vol.53. P.269-272.
- [13] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova. Phenolic profile of *Potentilla anserina* L. (Rosaceae) herb of Siberian origin and development of a rapid method for simultaneous determination of major phenolics in *P. anserina* pharmaceutical products by microcolumn RP-HPLC-UV. *Molecules*. **2015**. Vol.20. P.224-248.
- [14] N.I. Kashchenko, N.K. Chirikova, and D.N. Olennikov. Ellagitannins in Rosaceous plants from the flora of Sakha (Yakutia) Republic *Butlerov Communications*. **2014**. Vol.39. No.8. P.127-138. ROI: jbc-02/14-39-80-127
- [15] D.N. Olennikov, N.K. Chirikova. Phenolic compounds and cinnamamide from *Scutellaria scordiifolia*. *Chem. Nat. Comp.* **2013**. Vol.49. P.109-111.
- [16] <http://www.theplantlist.org/tpl1.1/record/kew-205377>.
- [17] Flora of Siberia. Ed. L.I. Malysheva. Vol.11: Pyrolaceae – Lamiaceae. *Novosibirsk: Nauka*. **1997**. 294p. (russian)
- [18] Synopsis of Yakutian flora: Сосудистые растения. *Novosibirsk: Nauka*. **2012**. 272p. (russian)
- [19] E.B. Pospelova, I.N. Pospelov. Flora of vascular plants of Taimyr and adjoining territories. Part 1. Annotated list of flora and its general analysis. *Moscow: KMK*. **2007**. 455p. (russian)
- [20] Thyme: The genus *Thymus*. Eds. E. Stahl-Biskup, F. Saez. *London, New York: Taylor&Francis*. **2002**. 516p.
- [21] S.V. Zhigzhitzhapova, A.N. Rabzhaeva, I.V. Zvontsov, L.D. Radnaeva. Chemical composition of essential oil of *Thymus baicalensis* Serg. growing in Transbaikalia. *Chemistry of Plant Raw Materials*. **2008**. No.1. P.73-76. (russian)
- [22] T.H. Oh, S.S. Kim, W.J. Yoon, J.Y. Kim, E.J. Yang, N.H. Lee, C.G. Hyun. Chemical composition and biological activities of Jeju *Thymus quinquecostatus* essential oils against *Propionibacterium* species inducing acne. *J. Gen. Appl. Microbiol.* **2009**. Vol.55. P.63-68.
- [23] S.V. Zhigzhitzhapova, A.N. Rabzhaeva, I.V. Zvontsov, L.D. Radnaeva. Comparative evaluation of componential profile of essential oils of *Thymus baicalensis* Serg. and *Thymus sibiricus* (Serg.) Klok. Et. Shost. *Vegetable resources*. **2008**. Vol.44. P.90-91. (russian)
- [24] O.R. Pereira, S.M. Cardoso. Overview on *Mentha* and *Thymus* polyphenols. *Curr. Anal. Chem.* **2013**. Vol.9. P.382-396.
- [25] N.I. Kashchenko and D.N. Olennikov. Spectrophotometric analysis of phenolic compounds medicinal calendulas (*Calendula officinalis* L.). Revision study of existing methods. *Butlerov Communications*. **2014**. Vol.37. No.1. P.146-155. ROI: jbc-02/14-37-1-146
- [26] P.D. Marin, R.J. Grayer, G.C. Kite, V. Matevski. External leaf flavonoids of *Thymus* species from Macedonia. *Biochem. Syst. Ecol.* **2003**. Vol.31. P.1291-1307.
- [27] A.B. Horwath, R.J. Grayer, D.M. Keith-Lucas, M.S.J. Simmonds. Chemical characterization of wild population of *Thymus* from different climatic regions in southeast Spain. *Biochem. Syst. Ecol.* **2008**. Vol.36. P.117-133.
- [28] A.F. Afonso, L.R. Pereira, R.T. Neto, A.M.S. Silva, S.M. Cardoso. Health-promoting effects of *Thymus herba-barona*, *Thymus pseudolanuginosus*, and *Thymus caespititius* decoctions. *Int. J. Mol. Sci.* **2017**. Vol.18. No 1879; doi:10.3390/ijms18091879.
- [29] L.M. Hernandez, F.A. Tomas-Barberan, F. Tomas-Lorente. Study of free flavone aglycones from some Iberian *Thymus* species. *Biochem. Syst. Ecol.* **1987**. Vol.15. P.61-67.