

Computation of vibrational spectra of *p*-nitrotoluene and 2,4,6-trinitrotoluene in the coordinates X_{δ}^0

© Alexander V. Belik,*[†] Erick G. Mischinsky, and Maxim Yu. Pavlichev

Faculty of Chemical Technology and Computing Chemistry. Chelyabinsk State University.

Kashyrinych Br. St., 129. Chelyabinsk, 454001. Russia. Phone: +7 (351) 799-70-66. E-mail: belik@csu.ru

*Supervising author; [†]Corresponding author

Keywords: *p*-nitrotoluene, 2,4,6-trinitrotoluene, the generalized force constants, coordinates X_{δ}^0 , calculations DFT, vibration frequencies (wavenumbers).

Abstract

In the framework of the density functional methods B3LYP 6-311++G(3df,3pd) first derived force field for *p*-nitrotoluene and 2,4,6-trinitrotoluene in the coordinates X_{δ}^0 . Determined generalized force coefficients for the relations in the coordinate X_{δ}^0 calculated frequency (wave number) of normal vibrations and carried out their assignment to certain types of vibrations.

Our results show that for *p*-nitrotoluene the most intense band in the vibrational spectrum has a value of 1374 cm⁻¹. Preferably in this vibration involves the atoms of the nitro group. It is a valence fluctuation of C-N bond and symmetric vibrations of the two bonds N-O, which are in opposite phase to C-N. Following the intensity band is mostly "owned" as nitrogroup. It matters 1577 cm⁻¹. This vibration involves the atoms of N and O. This is asymmetric vibrations of the two bonds N-O.

In the molecule of 2,4,6-trinitrotoluene has a characteristic intense bands involving nitro groups. The most intense band has a value of 1375 cm⁻¹. It could be attributed to the stretching vibrations of the nitro group. This is mainly the vibration of the bond C-N in position 4 and, in antiphase to that vibration, the symmetric vibrations of the two bonds N-O. The second intensity band with a value of 1604 cm⁻¹ corresponds to (mainly) the nitro groups in positions 2 and 6. This asymmetric vibrations of N-O in each of the nitro groups that have coordinated movement relative to each other. Wave number 1385 cm⁻¹ corresponds to the following intensity bond and is interpreted as asymmetric vibrations of nitro groups in positions 2 and 6 of the ring. This means that the C-N bond changes in the opposite phase each against the other. Inside each of the nitro groups N-O bonds change symmetrically.

The obtained generalized force coefficients of bonds in molecules of *p*-nitrotoluene and 2,4,6-trinitrotoluene in the coordinates X_{δ}^0 . So the generalized force coefficients for the relations C-N and N-O nitro group in the molecule, and *p*-nitrotoluene was found to be 6.6509 mdyn/Å and 12.7008 mdyn/Å (average) in the coordinates X_{δ}^0 , respectively. The generalized force coefficients for the relations C-N and N-O nitro group in the molecule 2,4,6-trinitrotoluene equal, respectively, 6.4682 mdyn/Å and 12.8707 mdyn/Å (middle).

References

- [1] S.S. Novikov, G.A. Shvehgeimer, V.V. Sevost'yanova, V.A. Shlyapochnikov. Chemistry of aliphatic and alicyclic nitrocompounds. *Moscow: Chemistry*. **1974**. 416p. (russian)
- [2] Chemical encyclopedic dictionary. Chief editor I.L. Knunyanc. *Moscow: Sov. Encyclopedia*. **1983**. 792p. (russian)
- [3] V.I. Sloveckii. The infrared absorption spectrum of aliphatic nitro compounds and their derivatives. *Successes of chemistry*. **1971**. Vol.XL. No.4. P.740-763. (russian)
- [4] A.V. Shastin, T.I. Godovikova, B.L. Korsunskii. Nitro-derivatives of 1,3,5-triazine. Synthesis and properties. *Successes of chemistry*. **2003**. Vol.72. No.3. P.311-320. (russian)
- [5] A.A. Lobanova, S.G. Il'yasov, G.V. Sakovich. Nitramide. *Successes of chemistry*. **2010**. Vol.79. No.9. P.894-905. (russian)
- [6] N.N. Mahova, A.S. Kulikov. Advances in the chemistry of monocyclic amino - and nitrofuraxans. *Successes of chemistry*. **2013**. Vol.82. No.11. P.1007-1033.
- [7] N.I. Sadova, L.V. Vil'kov. The molecular geometry of nitro-compounds. *Successes of chemistry*. **1982**. Vol.51. No.1. P.153-184.

- [8] A.V. Belik, R.Z. Zaharyan, V.A. Shlyapochnikov. The electronic structure of the molecules of nitro compounds. *Russian Chem. Bull. Ser.chem.* **1976**. No.8. P.1714-1719. (russian)
- [9] A.V. Belik. Modern elements computational chemistry: monograph. *Chelyabinsk: Publ. Chelyab. State Univ.* **2013**. 161p. (Classical University). (russian)
- [10] A.V. Belik New aspects of computation chemistry: monograph. *Chelyabinsk: Polygraph – Master.* **2015**. 200p. (russian)
- [11] A.V. Belik, I.L. Komkova, V.A. Shlyapochnikov. Theoretical analysis of the vibrational spectrum of nitromethane. *Russian Chem. Bull. Ser.chem.* **1985**. No.6. P.1410-1413. (russian)
- [12] I.V. Tokmakov, V.A. Shlyapochnikov. Using the theory of hybrid functional of the electronic density (hybridDFT) for the calculation of vibrational spectra: the nitromethane. *Russian Chem. Bull. Ser.chem.* **1997**. No.12. P.2106 -2109. (russian)
- [13] E.I. Fedotova, and A.V. Belik. Nitrocompounds: structure, force coefficients in coordinates X_8^0 and the frequencies of normal vibrations of NO_2 -group. *Butlerov Communications.* **2012**. Vol.31. No.9. P.29-35. ROI: jbc-02/12-31-9-29
- [14] A.V. Belik. Computation of vibrational spectra of benzene and benzofuroxan molecules in coordinates X_8^0 with an estimation of a force fields in frameworks DFT. *Butlerov Communications.* **2016**. Vol.45. No.2. P.148-162. DOI: 10.37952/ROI-jbc-01/16-45-2-148
- [15] A.V. Belik. Computation of vibrational spectra of 5-methoxy-4-nitrobenzofuroxan in coordinates X_8^0 . *Butlerov Communications.* **2016**. Vol.47. No.9. P.85-94. DOI: 10.37952/ROI-jbc-01/16-47-9-85
- [16] V. Kon. Electronic structure of matter – wave functions and density functionals. *Advances in physical Sciences (Nobel lectures in chemistry – 1998).* **2002**. Vol.172. No.3. P.336-348. (russian)
- [17] A.A. Korlyukov, M.Yu. Antipin. The study of the structure of crystals of organic and Organoelement compounds using modern quantum chemical calculations within the framework of the density functional theory. *Successes of chemistry.* **2012**. Vol.81. No.2. P.105-129.
- [18] W. Koch, M.C. Holthausen. A Chemist's Guide to Density Functional Theory. 2nded. *Weinheim: Wiley-Vch.* **2001**. 293p.
- [19] E.V. Butyrskaya. Computational chemistry: basic theory and work with programs Gaussian and Gauss View. *Moscow: SOLON-PRESS.* **2011**. 224p. (russian)
- [20] Gaussian 09, Revision A.02. M.J. Frisch, G.W. Trucks, H.B. Schlegel [at el.]. *Wallingford: Gaussian, Inc.* **2009**.
- [21] Spectral Database for Organic Compounds, SDBS [Электронныйресурс]. S. Kinugasa, K. Tanabe, T. Tamura. Базаданных. National Institute of Advanced Industrial Science and Technology (AIST), Japan. Режимдоступа: http://sdb.sdb.aist.go.jp/sdb/cgi-bin/cre_index.cgi
- [22] L.S. Mayanc. Theory and calculation of molecular vibrations. *Moscow: Publ. of Acad. Science of USSR.* **1960**. 526p. (russian)
- [23] E. Vil'son, Dj. Deshius, L. Kross. The theory of vibration spectra of molecules. *Moscow: Publ. «Foreign literature».* **1960**. 358p. (russian)
- [24] S. Sivin. Vibrations of molecules and middlesquare amplitude. *Moscow: Mir.* **1971**. 487p. (russian)
- [25] L.S. Mayanc, G.B. Shaltuper. A new approach to the complete calculation of the vibrations of any molecule. *Dokl. USSR Academy of Sciences.* **1972**. No.206. P.657-660. (russian)
- [26] L.S. Mayants, G.B. Shaltuper. General methods of analysing molecular vibrations. *J. Mol. Struct.* **1975**. Vol.24. P.409-431.
- [27] A.V. Belik. Solution of a spectral problem for molecules NH_2NO_2 , CH_3NO_2 and N_2O_4 in coordinates X_8^0 . *Butlerov Communications.* **2014**. Vol.40. No.11. P.84-90. ROI: jbc-02/14-40-11-84