

Reactivity indexes of alkenes aminomethylation

© Albert Kh. Fattakhov,¹⁺ Ivan V. Vakulin,^{1*} Azamat R. Galiakhmetov,²
Elvira R. Latypova,¹ and Rifkat F. Talipov^{1*}

¹Department of Organic and Bioorganic Chemistry. Chemistry Faculty. Bashkir State University.
Zaki Validi St., 32. Ufa. 450074. Republic of Bashkortostan. Russia.
Phone: +7 (347) 229-97-29. Email: al_fatt@mail.ru

²Department of Chemistry Marquette University Wehr Chemistry Building P.O.
Box 1881. Milwaukee, WI-53201. Email: azamatne@mail.ru

*Supervising author; [†]Corresponding author

Keywords: alkenes, aminomethylation, B3LYP and MP2 methods, NBO-analysis, reactivity indexes.

Abstract

Aminomethylation of alkenes is one of the simple and convenient method for obtaining various types of amines. The reaction proceeds under mild conditions and allows to introduce in one step -CH₂-NH₂ fragment, which suitable for subsequent transformations. Despite the similarity of the mechanisms of the reactions of Mannich and Prinse, the possibility of involving alkenes in the aminomethylation reaction is still insufficiently studied. So far, the range of alkenes involved in this reaction is very limited, and the prospects for its expansion have not been theoretically worked out. We believe that an important tool in assessing the reactivity of alkenes and, consequently, determining their activity in the reaction of aminomethylation are indexes of reactivity. Therefore, the search for correct reactivity indexes of alkenes in the aminomethylation reaction is an actual problem.

Consideration of the generally accepted indexes of reactivity such as partial charge on the atom, the occupancy of the orbitals, the bond orders makes it impossible to form an adequate model describing the reactivity of alkenes in the aminomethylation reaction.

It was shown that only NBO analysis of the electronic structure of alkenes the results of the calculations in the approximation B3LYP/6-31G(d,p) and MP2/6-31G(d,p) can be obtained in the correct model of the activity of alkenes. Correct indexes of reactivity, correctly taking into account the features of the reaction mechanism and the structure of alkenes with double bonds, are obtained by consideration of the uneven distribution of the electron density of π-bond of the double bond alkenes. The proposed indexes are in good agreement with the literature data on reactivity of alkenes in the aminomethylation reaction and can be used to predict their reactivity.

References

- [1] A.Kh. Fattakhov, R.F. Talipov, A.F. Kayumova, I.S. Shepelevich. Perspektivy aminometilirovaniya alkenov kak ekologichnogo sposoba polucheniya aminov. *Bashkir University Bulletin*. **2016**. Vol.21. No.4. P.893-905. (russian)
- [2] I.S. Shepelevich, Ye.YU. Ptashko, I.V. Vakulin, V.P. Timofeyev, R.F. Talipov. Kvantovo-
khimicheskoye issledovaniye vozmozhnosti aminometilirovaniya alkenov. *Bashkir Chemical Journal*. **2007**. Vol.14. No.1. P.134-136. (russian)
- [3] R.F. Talipov, I.V. Vakulin, A.R. Galyakhmetov, E.R. Latypova, and G.R. Talypova. Reactivity indices of alkenes in amino-methylation reactions. *Butlerov Communications*. **2014**. Vol.39. No.7. P.47-52. ROI: jbc-02/14-39-7-47
- [4] G. Henonion, C. Price, V. Wolff. The reaction of formaldehyde and secondary amines with some olefins. *J. Am. Chem. Soc.* **1955**. Vol.77. No.17. P.4633-4636.
- [5] C.J. Schmidle, R.C. Mansfield. The reaction of secondary amines, formaldehyde and olefin. *J. Am. Chem. Soc.* **1955**. Vol.77. No.17. P.4636-4638.

- [6] C.J. Schmidle, R.C. Mansfield. The Aminomethylation of Olefins. II. A New Synthesis of 1-Alkyl-4-aryl-4-piperidinols. *J. Am. Chem. Soc.* **1955**. Vol.77. No.21. P.5698-5700.
- [7] T. Cohen, A.J. Onopchenko. Competing hydride transfer and ene reactions in the aminoalkylation of 1-olefins with N,N-dimethylmethyleniminium ions. A literature correction. *Org. Chem.* **1983**. Vol.48. No.24. P.4531-4537.
- [8] <http://classic.chem.msu.su/gran/gamess/index.html>
- [9] Weinhold F "Natural Bond Orbital Methods" in Schleyer Pv.R., Allinger N.L., Clark T., Gasteiger J., Kollman P.A., Schaefer H.F., Schreiner P.R. (Eds.) Encyclopedia of Computational Chemistry. John Wiley & Sons, Chichester, UK, 3. **1998**. P.1792-1811.
- [10] Glendening E.D., Landis C.R., Weinhold F. Natural bond orbital methods. *Wiley Interdisciplinary Reviews-Computational Molecular Science*. **2012**. Vol.2. P.1-43.
- [11] H-Z. Li, Y-C. Wang, Z-Y. Geng, Q-L. Zhang, Q-Y. Wang, Si Y-B. Yu-Bing. The theoretical investigation on gas-phase chemistry of YNH^+ with propene. *J. of Mol. Struct: THEOCHEM*. **2008**. Vol.866. P.5-10.
- [12] A.S. Khamidullina, I.V. Vakulin, R.F. Talipov, I.S. Shepelevich. Structure effects of the protonated lincomycin molecule on the mechanism of its complexation with organic compounds. *J. of Struct. Chem.* **2005**. Vol.46. No.6. P.985-990.
- [13] K.Ya. Burshteyn, P.P. Shorygin. Quantum chemical calculations in organic chemistry and molecular spectroscopy. Moscow: Nauka. **1989**. P.104. (russian)
- [14] A.E. Reed, R.B. Weinstock, F.J. Weinhold. Natural Population Analysis. *Chem. Phys.* **1985**. Vol.83. P.735-746.
- [15] J.E. Carpenter, F. Weinhold. Analysis of the geometry of the hydroxymethyl radical by the "different hybrids for different spins" natural bond orbital procedure. *J. of Mol. Struct: THEOCHEM*. **1988**. Vol.169. P.41-62.
- [16] M. Arivazhagan, R. Kavitha. Molecular structure, vibrational spectroscopic, NBO, HOMO-LUMO and Mulliken analysis of 4-methyl-3-nitro benzyl chloride. *J. of Mol. Struct.* **2012**. Vol.1011. P.111-120.
- [17] S.G. Semenov, E.S. Apostolova, S.M. Shevchenko. The structure of 1,4'-dihydroxybenzyl cation. *J. of Mol. Struct: THEOCHEM*. **1991**. Vol.251. P.389-394.
- [18] H. Junjappa, H. Ila, C.V. Asokan. α -Oxoketene-S,S-, N,S- and N,N-acetals: Versatile intermediates in organic synthesis. *Tetrahedron*. **1990**. Vol.46. No.16. P.5423-5506.
- [19] D.H.R. Barton, R.V. Stick, R. Subramanian. *J. Chem. Soc., Perkin Trans. I*. **1976**. Vol.19. P.2112-2116.
- [20] P. Geerlings, A.M. Vos, R.A. Schoonheydt. A computational and conceptual DFT approach to the kinetics of acid zeolite catalyzed electrophilic aromatic substitution reactions. *J. of Mol. Struct: THEOCHEM*. **2006**. Vol.762. No.1. P.69-78.
- [21] M.A. Battiste, J.M. Coxon, G.W. Simpson, P.J. Steel, A.J. Jones. Orbital control of stereochemistry in acid-catalysed addition reactions of endo-tricyclo[3.2.1.0^{2,4}]oct-6-ene. *Tetrahedron*. **1984**. Vol.40. No.16. P.3137-3144.
- [22] M. Ishida, M. Sakamoto, H. Hattori, M. Shimizu, S. Inagaki. Orbital control of π -facial selectivity in Diels-Alder reactions of cyclopentadienes having C(=O)YR substituents at the 5-positions. *Tetrahedron Lett.* **2001**. Vol.42. No.20. P.3471-3474.
- [23] E.K. Kimanani, J. Lavigne. Bioanalytical calibration curves: variability of optimal powers between and within analytical methods. *J. Pharm. Biomed. Anal.* **1998**. Vol.16. No.6. P.1107-1115.
- [24] E.K. Kimanani. Bioanalytical calibration curves: proposal for statistical criteria. *J. Pharm. Biomed. Anal.* **1998**. Vol.16. No.6. P.1117-1124.