

## Agroecological assessment of soil structure

© Sergey L. Belopukhov,<sup>1\*</sup> Vitaly I. Savich,<sup>1</sup> Viktor V Gukalov,<sup>2</sup>  
Nuruly M. Saduakasov,<sup>1</sup> Kristina S. Los,<sup>1</sup> and Elena M. Ilyushkova<sup>1</sup>

<sup>1</sup>Russian Timiryazev State Agrarian University. Timiryazevskaya St., 49. Moscow, 127550.

Russia. Phone: +7 (499) 976-28-62. E-mail: SBelopuhov@rgau-msha.ru

<sup>2</sup>North-Caucasian Experimental Station. Krasnodarski Region. Russia.

\*Supervising author; †Corresponding author

**Keywords:** structure of soils, pedes, kutans, fertility of soils.

### Abstract

The formation of the soil structure is determined by the soil formation factors, by the soil-forming processes, but at the same time, the structure formed in the soil affects the physico-chemical and microbiological processes in the soil profile and in the catheter, and determines the fertility of soils.

In the studies carried out, the chemical composition of the fractions of a lumpy structure of different sizes is distinguished. In the chernozems, 10-3 mm fractions contained (in the extract  $\text{CH}_3\text{COOHN}_4$ ) less than Ca –  $23.9 \pm 2.1 (\cdot 10^3)$  mg/l, Mg –  $9.5 \pm 0.2 (\cdot 10^3)$ ; K –  $102.2 \pm 3.3$ , and the fractions of 2-0.25 mm, respectively, Ca –  $29.9 \pm 1.9 (\cdot 10^3)$ ; Mg –  $10.8 \pm 0.7 (\cdot 10^3)$ ; K –  $120.9 \pm 3.6$ .

The outer and inner layers of prismatic structural separations of sod-podzolic soil differed in the content of water-soluble forms of calcium and magnesium. In poorly cultivated soil, Mg, Ca, respectively, in the outer layer 3.2; 2.6 mg/l; in the inner layer 4.2; 3.5 mg/l. In well-cultivated soil in the outer layer, Mg is 8.6, Ca is 7.6; K – 1.2 mg/l; in the inner case, respectively, 3.9; 3.8 and 0.9 mg/l. The planes and edges of the structural units differed in biological activity.

The introduction of stubble residues of plants and organic fertilizers led to an increase in the coefficient of soil structure. When 3 g of crop residues per 100 g of soils were introduced into the sod-podzolic soil, the aggregate content of more than 3 mm increased from 0.11% to 9.39%. The introduction of organic-mineral compost in the field to chernozems led to an increase in the structural coefficient from 2.5 to 3.0.

### References

- [1] R.A. Gadzhiagaeva Spatial differentiation of the state and behavior of  $^{90}\text{Sr}$  at the aggregate level of soils, Moscow: *RGAU-MAHA, Abstract of thesis. Cand. diss.* **2015**. 24p. (russian)
- [2] V.V. Gukalov The influence of complex organo-mineral composts on the properties and processes in the soil-plant system on ordinary chernozem, the development and productivity of corn sowing, *Author's Abstract of thesis. Cand. Diss., Moscow, VNIIA.* **2015**. 19p. (russian)
- [3] V.N. Gukalov, V.I. Savich, I.S. Beliuchenko. Information-energy assessment of the state of heavy metals in landscape components. Moscow: *RGAU-MSHA, VNIIA.* **2015**. 400p. (russian)
- [4] E.V. Dubovik. Interrelation of the humus and structural state in eroded chernozems and slope agrarian landscapes of the Central Chernozem Region, *Author's abstract. Doct. Diss. Moscow: RGAU-MAHA.* **2017**. 48p. (russian)
- [5] A.G. Zamaraev, V.I. Savich, V.G. Sychev and others. Energy-exchange in the link of the field crop rotation. Part 2. Moscow: *RGAU-MAKHA.* **2005**. 336p. (russian)
- [6] Yu.M. Klovskaia. Root entry of  $^{137}\text{Cs}$  into plants from aggregated soils at the initial stages of radionuclide contamination and over time, *Abstract. Cand. Diss. Moscow: RGAU-MAKSA.* **2015**. 24p. (russian)
- [7] A.R. Rahmatullina. Composition and properties of clayey material of sod-podzolic soils of different degree of cultivation and their role in the formation of structural units, *Author's Abstract of. Cand. Diss. St.-Petersburg.* **1992**. 21p. (russian)
- [8] V.I. Savich, S.L. Belopukhov, M.D. Alifirov, G.N. Kushnir, I.G. Shaykhiev. Agroecological aspects in the allocation of humates from plant biomass and organic fertilizers. 1. Effect of humates on soil properties and biotests. *Bulletin of Kazan University of Technology.* **2015**. Vol.18. No.20. P.238-241. (russian)

**Full Paper** \_\_\_\_\_ S.L. Belopukhov, V.I. Savich, V.V. Gukalov, N.M. Saduakasov, K.S. Los, and E.M. Ilyushkova

- [9] T.V. Shnee, S.E. Starykh, T.A. Fedorova and others. Changes in the physical and chemical properties of soil colloids depending on the ionic composition of the soil absorbing complex. *Fertility*. **2014**. No.3(78). P.33-35. (russian)
- [10] V.I. Savich, V.S. Krutilina, D.N. Egorov, A.D. Kashansky. Use of computer diagnostics for an objective characterization of soil color. *Izvestiya TSHA*. **2004**. No.4. P.38-51. (russian)
- [11] V.I. Savich, V.D. Naumov. Local processes of soil-forming processes as a factor of correction of fertility models of soils. *International Agricultural Journal*. **2017**. No.1. P.49-53. (russian)
- [12] V.I. Savich, M. Mazirov, B.A. Borisov. Assessment of the optimal soil properties and the lack of nutrients for plants using methods based on the principles of feedback. *International Agriculture*. **2017**. No.4. P.48-50. (russian)
- [13] I.N. Stepanov. Forms in the world of soils. *Moscow: Nauka*. **1986**. 192p. (russian)
- [14] A.D. Fokin, S.P. Torshin, Yu.M. Bebneva and others. The intake of <sup>137</sup>Cs, <sup>90</sup>Sr from the surface of soil aggregates and from their intra-pedal space. *Pochvovedenie*. **2014**. No.12. P.1416-1425. (russian)
- [15] R.F. Khasanova. Agroecological analysis of the structural state and optimization of the properties of the chernozems of the Transural region in phytomelioration. *Author's abstract of. Doct. Diss.*, Ufa. **2016**. 42p. (russian)
- [16] V.I. Savich, S.L. Belopukhov, V.V. Gukalov. Complexing ability of decomposition products of plant litter and its effect on the mobility of polyvalent cations. *Bulletin of Kazan University of Technology*. **2015**. Vol.18. No.12. P.185-188. (russian)