

## Analysis of the mechanism of radical chain oxidation of ethylbenzene in the presence of additive of *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine

© **Gulnaz M. Sharipova,<sup>1+</sup> Irina V. Safarova,<sup>1</sup> Anatoly Ya. Gerchikov,<sup>1\*+</sup> Ildus Sh. Nasyrov,<sup>2</sup> and Rozalia D. Bulyakova<sup>1</sup>**

<sup>1</sup>*Department of Physical Chemistry and Chemical Ecology. Bashkir State University.*

*Z. Validi St., 32. Ufa, 450076. Republic of Bashkortostan. Russia.*

*Phone: +7 (347) 273-67-27. Fax: +7 (347) 273-67-01.*

*E-mail: gerchikov@inbox.ru,<sup>1</sup> gulnaz-sharipova@list.ru*

<sup>2</sup>*Open Joint Stock Company "SyntezCauchuk". Tekhnicheskaya St., 14. Sterlitamak, 453110. Russia.*

\*Supervising author; <sup>+</sup>Corresponding author

**Keywords:** *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine, inhibitor, mathematical model, kinetics.

### Abstract

A primary consideration of modern production in chemistry, medicine and the food industry is preventing unwanted oxidation processes. Currently, the quantitative characteristic of the antioxidant activity of potential antioxidant is the value of the inhibition rate constants, whose definition is available by known methods of chemical kinetics. To select an effective inhibitor of oxidative processes, it is necessary to define the optimum conditions of its validity and maximum braking, that it can provide in the process that becomes possible only after establishing the mechanism of its action. It is not sufficient for the application of methods of physical-chemical experiment for an explanation of the mechanism of action of antioxidants, because the process that inhibited by oxidation includes a large number of stages involving intermediate particles-atoms and radicals concentration and lifetime of them is extremely small. A necessary component that allows us to solve this complex problem is mathematical modeling of mechanism of action of compounds based on the available experimental results.

Therefore, the aim of this work was the study of the mechanism of antioxidant action of *N*-2-ethylhexyl-*N'*-phenyl-*p*-phenylenediamine (Novantoks, 8-PFDA) reactions initiated by the oxidation of ethylbenzene by using mathematical modeling. To achieve this goal, was used the program complex "HimKinOptima" and the obtained previously experimental data. By solving the inverse problem of chemical kinetics was restored everything, including previously unknown, key rate constants of the elementary stages. By solving the direct problem of chemical kinetics was received full kinetic painting of all the reaction participants, including unstable intermediates. It is shown that the presence of impurities of alcohol in the starter compound leads to an increase in its antiradical action at the expense of regeneration reaction inhibitor in the act of truncating the chain with the participation of peroxy radicals.

The reliability of research results obtained in theory ensured the correctness of computer modeling methods used in accordance with the established objectives and ensures a satisfactory settlement and coincidence experimental significance of quantitative parameters.

### Литература

- [1] M. Kavun, Yu.V. Vinokurov, A.I. Efremov, I.G. Ahmetov, N.M. Evtishina, V.N. Borisenko, E.A. Tulchinskiy. Novantoks 8 PFDA-best antioxidant for rubber refills-3. *The world of tire*. **2006**. No.8. P.14-22. (russian)
- [2] N.I. Koltsov, N.F. Ushmarin, L.G. Rogozhina, S.A. Issakova, A.V. Jarutkina, A.Y. Plehanova, and M.V. Kuzmin. Research of influence of technological additives on properties of rubbers on the basis of BNR new generation. Part 4. Powder stabilizers on a basis novantox 8 PFDA. *Butlerov Communications*. **2010**. Vol.22. No.10. P.42-50. ROI: jbc-02/10-22-10-42
- [3] M.V. Tikhonova, D.F. Maskov, S.I. Spivak, I.M. Gubaidullin. Program complex "HimKinOptima" for mathematical modeling and optimization based on kinetics of chemical reactions using parallel

- ANALYSIS OF THE MECHANISM OF RADICAL CHAIN OXIDATION OF ETHYLBENZENE IN THE PRESENCE... 102-106  
computing and database: certificate of registration of an electronic resource. INIPI RAO OPHERNiO. No  
19247; date of reg. 30.05.2013. (russian)
- [4] M.V. Tikhonova, G.G. Garifullina, A.Ya. Gerchikov, S.I. Spivak. The kinetic model of n-decane oxidation in the presence of inhibitory composition. *Int. J. Chem. Kinet.* **2014**. Vol.46. No.4. P.220.
- [5] A.Ya. Gerchikov, G.R. Ahatova, G.M. Sharipova, A.G. Mustafin, M.V. Sahibgareeva, S.I. Spivak. Research of mechanism of inhibited oxidation of 1,4-dioxane by method of mathematical modeling. *Kinetics and catalysis.* **2015**. Vol.56. No.3. P.301-304. (russian)
- [6] A.Ya. Gerchikov, G.M. Sharipova, G.R. Ahatova, A.G. Mustafin, M.V. Sahibgareeva, S.I. Spivak. Kinetics and mechanism of reaction of uracil with hydrogen peroxide in aqueous solutions. Mathematical model of reaction. *Kinetics and catalysis.* **2015**. Vol.56. No.5. P.571-576. (russian)
- [7] M.V. Tikhonova, G.G. Garifullina, A.Ya. Gerchikov, S.I. Spivak. Mathematical modeling of radical chain oxidation of organic compounds. *Kinetics and catalysis.* **2013**. Vol.54. No.4. P.429-433. (russian)
- [8] A.Ya. Gerchikov, G.M. Sharipova, I.V. Safarova, M.V. Sahibgareeva, S.I. Spivak. Reactivity and mechanism of action of fullerene C<sub>60</sub> as radical chain oxidation inhibitor 1,4-dioxane. *Physical chemistry journal.* **2017**. Vol.96. No.6. P.957-962. (russian)
- [9] G.M. Sharipova, I.V. Safarova, A.Ya. Gerchikov, I.Sh. Nasyrov, and G.N. Musina. N-2-ethylhexyl-N'-phenyl-p-phenylenediamine as inhibitor of radical chain oxidation of ethylbenzene. *Butlerov Communications.* **2017**. Vol.50. No.6. P.126-131. DOI: 10.37952/ROI-jbc-01/17-50-6-126
- [10] Yu.V. Vinokurov, S.M. Kavun, N.I. Koltsov. Development of process of receiving N-2-ethylhexyl-N'-phenyl-p-phenylenediamines on the basis of a research of features of his synthesis in laboratory and industrial condition. *Bulletin of the Kazan Technological University.* **2011**. No.15. P.48-56. (russian)
- [11] E.T. Denisov, T.G. Denisova. Oxidation of alcohols with two types of peroxy radicals. *Petrochemistry.* **2006**. Vol.46. No.5. P.333-340. (russian)
- [12] E.T. Denisov. Liquid-Phase Reaction Rate Constants. *New York: Plenum.* **1974**. P.1-711.
- [13] N.M. Emanuel, E.T. Denisov, Z.K. Mayzus. Chain reactions of oxidation of hydrocarbons in a liquid phase. *Moscow: Nauka.* **1965**. (russian)