

Noncatalytic oxidation of sulfur dioxide

© Aigul I. Tuktarova,¹⁺ Linara R. Baraeva,^{1*} Guzelya I. Sabahova,²
Alsu A. Yusupova,¹ and Rezida T. Akhmetova¹

¹Technology of Inorganic Substances and Materials Division. Institute of Petroleum, Chemistry and Nanotechnologies. Kazan National Research Technological University. K. Marx St., 68. Kazan, 420015. Republic of Tatarstan. Russia. Phone: +7 (843) 238-56-94.

E-mail: office@kstu.ru, baraeva.linara@yandex.ru

²Enhanced Oil Recovery Department. Tatar Oil Research and Design Institute of Tatneft Company (TatNIPIneft). M. Jalil St., 32. Bugulma, 423230. Republic of Tatarstan. Russia. Phone: +7 (855) 947-85-55. E-mail: sabahova.guzel@yandex.ru

*Supervising author; +Corresponding author

Keywords: sulfur dioxide, sulfuric acid, sulfur trioxide, oxidation, activation energy, quantum chemical calculations.

Abstract

The theoretical research of process of oxidation dioxide sulfur without catalyst with use of the quantum and chemical Priroda 6 program is conducted. This program is an available software package for scientific and study, she doesn't demand big time expenditure and allows to receive high precision of calculations, and there is also an opportunity to carry out parallel calculations. The ChemCraft program visualizer was used for process modeling. During the sulfur dioxide oxidation process mechanism research for a singletny state it is revealed that this process is difficult and multistage, during reaction formation of several products is possible. At the first stage of the studied process two molecules SO₃ separate, further there is a regrouping to formation of the SO₅ and SO complex. Then the way of process is divided into two in parallel the proceeding stages. One of stages leads to formation, most likely, of the final products of SO₃, SO and O₂. Another in parallel the proceeding stage leads to formation of by-products of SO₄ and SO₂. The most power-intensive stage is the second, E act = 232 kJ/mol. On the obtained data the power chart of a way of reaction of oxidation of dioxide of sulfur is constructed. For all sites of process thermodynamic characteristics are calculated, and for stages values of energy of activation and thermal effects are found. All stages, except the second, go spontaneously with low energy of activation, the third stage is surmountable in temperature conditions of synthesis. The most part of an activation barrier of homogeneous oxidation of dioxide of sulfur is caused by energy expense for a rupture of communication of S-O and transition of one atom of oxygen to the molecule SO₄ with formation of the SO₅ complex. The obtained data, namely the second energy of activation, prove need of carrying out process in the presence of the catalyst. At selection of catalysts of process of oxidation it is necessary to consider probability of a molecule of dioxide of sulfur to react with oxygen atoms, connected with the catalyst less strongly, than with each other in the molecule O₂, for overcoming a power barrier.

References

- [1] Information on contact oxidation of sulfur dioxide [electronic resource] https://studopedia.ru/7_45597_kontaktnoe-okislenie-okside-seri-IV.html (russian)
- [2] L.R. Baraeva, R.T. Akhmetova, A.I. Khatsrinov, G.I. Sabakhova. The use of the quantum-chemical program "Priroda" in the educational process of the TISM Department of the KNRTU. *Bulletin of Kazan Technological University*. **2013**. No.24. P.261-263. (russian)
- [3] K.S. Krasnov. Molecular constants of inorganic compounds: Proc. Allowance. *Leningrad: Chemistry*. **1979**. 448p. (russian)
- [4] L.R. Baraeva, G.I. Sabahova, and R.T. Akhmetova. Quantum-chemical modeling of process iron and zinc sulfides synthesis from their chlorides. *Butlerov Communications*. **2015**. Vol.41. No.2. P.141-146. DOI: 10.37952/ROI-jbc-01/15-41-2-141