

Investigation of molecular of clusters of petroleum asphaltenes

© Svetlana A. Shutkova,¹⁺ Mikhail Yu. Dolomatov,^{2,3*} Rauf Z. Bakhtizin,²
 Ildar R. Kharudinov,⁴ Milana M. Dolmatova,² and Zagir Z. Ishniyazov²

¹ Department of Power Engineering and Physical. Faculty of Energy Engineering.

Bashkir State Agrarian University. 50th anniversary of October, 34. Ufa. 450001. Bashkortostan Republic.

Russia. Phone: +7 (347) 228-52-00. E-mail: Svetlana-Shutkova@yandex.ru

² Department of Physical Electronics and Nanophysics. Physical and technical Institute.

Bashkir State University. Z. Validi St., 32. Ufa. 450001. Bashkortostan Republic. Russia.

Phone: +7 (347) 229-96-47. E-mail: dolomatov@mail.ru

³ Department of Technology of Oil and Gas. Faculty of Technology. Ufa State oil Technical University. Kosmonavtov St., 1. Ufa. 450062. The Republic Of Bashkortostan. Russia.

Phone: +7 (347) 243-15-35. E-mail: dolomatov@mail.ru

⁴ GUP Institute of Oil Refining Rep. Bashkortostan. Initiative St., 12. Ufa, 450065. Russia.

Phone/fax: +7 (347) 242-25-11, +7 (347) 242-24-73. E-mail: inhp@inhp.ru

*Supervising author; ⁺Corresponding author

Keywords: oil asphaltenes, nanoclusters, molecular mechanics method, ionization potential, electron affinity, electronic phenomenological spectroscopy.

Abstract

A study of the electronic and chemical structure of nanoclusters, consisting of molecular fragments of oil asphaltenes – "continental" type. The objects of study are asphaltenes residue of thermal cracking and tar Western-Siberian oil and asphaltenes of West Siberian oil. For experiments on the isolation and separation of asphaltenes was used the method of I.R. Khairudinov. The obtained electronic absorption spectra of solutions of asphaltenes West Siberian oil in the visible and UV region (280-780 nm). Processed spectra to a computer and calculating the effective potential of ionization and electron affinity of the studied objects by means of electronic phenomenological spectroscopy. The calculated values of the effective ionization potentials are in the range from 5.56 to 5.86 eV, electron affinity, from 1.68 eV to 1.91 eV. Physico-chemical properties of petroleum asphaltenes determined by correlation of range-property: value srednetsenovoj molecular weight (from 2437 to 3884 e.m.), energy of activation of viscous flow (162.4 to 272.4 kJ/mol), concentration of paramagnetic centers of the carbon (from 145.6 to 273.4 10¹⁸ spin/cm³). The calculated electronic structure of molecular nanoparticles of petroleum asphaltenes by the method DFT/B3LYP with basis set 6-31+G*.

According to the findings of the adiabatic first ionization potentials are in the range of 6.07 to 7.39 eV, the electron affinity is from 0.88 to 1.22 eV. The settlement data confirm the hypothesis that increased donor-acceptor abilities of asphaltic substances. The study of the structural characteristics of the nanoclusters petroleum asphaltenes using the method of molecular mechanics. The dihedral angle between the virtual planes of the alkyl ring and the plane of the ring is in the range from 0 to 108 1580. Thus the alkyl groups substituting hydrogen in the aromatic rings on the periphery of the substantially non-planar plane naphthenoaromatic rings. Confirmed deplanement structure naphthenoaromatic fragments of clusters. The dihedral angle between the plane of naphthenic rings and the plane of the aromatic rings in the structures of the nanoparticles has different values, being in the range from 161⁰ to 168⁰. The distance h between the virtual planes naphthenoaromatic fragments of the nanocluster is in the range from 3.4 to 3.7 Å. The possibility of the formation of nanoclusters of petroleum asphaltenes, consisting of naphtene aromatic plates. The calculation results showed that the values of intermolecular interaction of molecular fragments are in the range from 7 to 226 kJ/mol. The results of the rheological studies indicate high values of activation energy of viscous flow.

References

[1] D. Speight. Analysis of petroleum. Handbook. Moscow: Profession. 2010. 480p. (russian)

- [2] C.D. Sharma, S.D. Bhagat, S.Z. Erhan. Maltenes and Asphaltenes of Petroleum Vacuum Residues: Physico-Chemical Characterization. *Petroleum Science and Technology*. **2007**. No.25. P.93-104.
- [3] K. Rajagopal, S.M. Silva. An experimental study of asphaltene particle sizes in n-heptane-toluene mixtures by light scattering. *Brazilian Journal of Chemical Engineering*. **2004**. Vol.21. No.4. P.601.
- [4] D. Merino-Garcia, J.M. Shaw, H. Carrier, H.W. Yarranton, L. Goual. Petrophase 2009 Panel Discussion on Standardization of Petroleum Fractions. *Energy Fuels*. **2010**. Vol.24. No.4. P.2175-2177.
- [5] P.M. Spiecker, K.L. Gawrys, C.B. Trail, P.K. Kilpatrick, P.M. Spiecker. Effects of petroleum resins on asphaltene aggregation and water-in-oil emulsion formation. *Colloids and Surfaces A: Physicochem. Eng. Aspects*. **2003**. Vol.220. No.1. P.9-27.
- [6] S. Mitra-Kirtley, O.C. Mullins, J. Van Elp, S.J. George, J. Chen, S.P. Cramer. Determination of the nitrogen chemical structures in petroleum asphaltenes using XANES spectroscopy. *J. Am. Chem. Soc.* **1993**. Vol.115. No.1. P.252-258.
- [7] A.R. Hortal, B. Martinez-Haya, M.D. Lobato, J.M. Pedrosa, S. Lago. On the determination of molecular weight distributions of asphaltenes and their aggregates in laser desorption ionization experiments. *J. Mass Spectrom.* **2006**. Vol.41. No.7. P.960-968.
- [8] N.V. Lisitza, D.E. Freed, P.N. Sen, Y.Q. Song. Study of Asphaltene Nanoaggregation by Nuclear Magnetic Resonance (NMR). *Energy Fuels*. **2009**. Vol.23. No.3. P.1189-1193.
- [9] S.S. Betancourt, G.D. Ventura, A.E. Pomerantz, O. Vilorio, F.X. Dubost, J. Zuo, G. Monson, D. Bustamante, J.M. Purcell, R.K. Nelson, R.P. Rodgers, C.M. Reddy, A.G. Marshall, O.C. Mullins. Nanoaggregates of asphaltenes in a reservoir crude oil and reservoir connectivity. *Energy Fuels*. **2009**. Vol.23. No.3. P.1178-1188.
- [10] M.R. Gray, R.R. Tykwinski, J.M. Stryker, X. Tan. Supramolecular Assembly Model for Aggregation of Petroleum Asphaltenes. *Energy Fuels*. **2011**. Vol.25. No.7. P.3125.
- [11] T.F. Yen, J.G. Erdman, S.S. Pollack. Investigation of the Structure of Petroleum Asphaltenes by X-Ray Diffraction. *Anal. Chem.* **1961**. Vol.33. No.11. P.1587.
- [12] S.R. Bagheri, A. Bazyleva, M.R. Gray, W.C. McCaffrey, J.M. Shaw. Observation of Liquid Crystals in Heavy Petroleum Fractions. *Energy & Fuels*. **2010**. Vol.24. No.8. P.4327.
- [13] H. Groenzin, O.C. Mullins. Molecular Size and Structure of Asphaltenes from Various Sources. *Energy Fuels*. **2000**. Vol.14. No.3. P.677.
- [14] H. Sabbah, A.L. Morrow, A.E. Pomerantz, R.N. Zare. Evidence for Island Structures as the Dominant Architecture of Asphaltenes. *Energy Fuels*. **2011**. Vol.25. No.4. P.1597-1604.
- [15] D.E. Dmitriev, A.K. Golovko. Modeling the molecular structure of petroleum resins and asphaltenes and their thermodynamic stability calculations. *Chemistry for Sustainable Development*. **2010**. No.18. P.171-180.
- [16] J. Murgich. Physical and chemical properties of disperse systems and oil and gas technologies. *Izhevsk*. **2007**. 580 p. (russian)
- [17] J. Murgich, J.A. Abanero, O.P. Strausz. Molecular Recognition in Aggregates Formed by Asphaltene and Resin Molecules from the Athabasca Oil Sand. *Energy Fuels*. **1999**. Vol.13. No.2. P.278-286.
- [18] Yu. Ganeeva, T.N. Yusupova, G.V. Romanov. Structural organization of the asphaltenes. *Reports of the Academy of Sciences*. **2009**. Vol.426. No.5. P.629-631. (russian)
- [19] S.A. Sozinov, L.V. Sotnikova, A.N. Popova, and R.P. Kolmikov. Study of the film of n-hexane insoluble asphaltene produced from coal-tar pitch. *Butlerov Communications*. **2017**. Vol.51. No.7. P.75-80. ROI: jbc-02/17-51-7-75
- [20] M.Yu. Dolomatov, S.A. Shutkov, S.V. Dezortsev. Structure of molecular nanoparticles of petroleum asphaltenes. *Journ. Structur. Chem.* **2012**. Vol. 53. No.3. P.563-567. (russian)
- [21] M.Yu. Dolomatov, S.A. Shutkov, S.V. Dozortsev. Structural and Chemical Characteristics of Model Molecular Fragments of Petroleum Asphaltenes. *Petroleum Chemistry*. **2012**. Vol.52. No.4. P.299-303. (russian)
- [22] M.Yu. Dolomatov, S.A. Shutkova. Study of a Supramolecular Structure of Continental Type Petroleum Asphaltenes. *Journ. Structur. Chem.* **2017**. Vol.58. No.7. P.1311-1316. (russian)
- [23] M.Y. Dolmatov, A.B. Marushkin, R.N. Gimaev, M.M. Seliverstov. Thermodynamics of formation of supramolecular structure of asphaltenes. *Solid fuel chemistry*. **1989**. No.6. P.83-864. (russian)
- [24] I.R. Khairudinov, F.M. Sultanov, E.G. Telyashev. Modern processes of solvent deasphalting of oil residues. Ufa: Publishing house of SUP IPCP RB. *Series "Oil refining Library"*. **2011**. 208p. (russian)
- [25] M.Yu. Dolomatov, G.R. Mukaeva. A method for determining the potential of ionization and electron affinity of atoms and molecules by electron spectroscopy. *Journal of appl. Spectroscopies*. **1992**. Vol.56. No.4. P.570-575. (russian)

- Full Paper** S.A. Shutkova, M.Yu. Dolomatov, R.Z. Bakhtizin, I.R. Kharudinov, M.M. Dolmatova, and Z.Z. Ishniyazov
- [26] G.R. Mukaeva, M.Y. Dolmatov. Spectroscopic control of the properties of organic substances and materials property correlations is the absorption coefficient. *Journal of appl. spectroscopies*. **1998**. Vol.65. No.3. P.438-440. (russian)
- [27] F. Jensen. Introduction to Computational Chemistry. *USA, New York: John Wiley&Sons*. **2007**. 599p.
- [28] G.G. Valyavin, M.Yu. Dolomatov, A.I. Ilyasov, N.F. Yurchenko. Physico-chemical peculiarities of thermolysis of complex hydrocarbon systems. Experiment. Theory. Technology. *St.-Petersburg: Subsoil*. **2017**. 352p. (russian)