

## The role of phosphatidylethanolamine in the biogenesis of alkaline phosphatase in *Escherichia coli*

© Nataliia A. Bondareva,\*<sup>+</sup> Viktor V. Golovanov, and Alpha S. Gilmuddinova  
Chair of Organic, Bioorganic and Medicinal Chemistry. Samara National Research University.  
Akad. Pavlova, St., 1. Samara, 443011. Samara Region. Russia.  
Phone: +7 (846) 334-54-59. E-mail: nnkk86@mail.ru

\*Supervising author; <sup>+</sup>Corresponding author

**Keywords:** phosphatidylethanolamine, biogenesis, alkaline phosphatase, *Escherichia coli*.

### Abstract

Protein secretion, as is known, is determined by the primary structure of the protein and its specific domains, which contain information on the interaction with the membrane and components of the secretory apparatus. However, the structural principles of interaction between these components and the secreted protein are not fully established. This is especially true for the interaction of topogenic protein sequences with membrane phospholipids and, in particular, with phosphatidylethanolamine. Previously was shown that the signal peptide interacts with anionic phospholipids. However, the nature of this interaction *in vivo* has not been fully established. It is not clear whether a direct electrostatic interaction occurs between the N-terminal region of the SP (signal peptide) and anionic phospholipids, or the protein components of the secretory apparatus are involved. It remains an open question whether phosphatidylethanolamine (PEA) participates in the interaction of the signal peptide with membranes. Alkaline phosphatase is a typical secreted protein of *E. coli*, localized in the periplasm.

In our study described in this paper, the need for *E. coli* in PEA for the secretion of alkaline phosphatase *in vivo* was assessed and the relationship between this requirement and the involvement of PEA in the formation of the non-bilayer structure was identified.

Despite the fact that the protein components of the secretory apparatus are sufficiently well studied and characterized, it is impossible to understand the molecular mechanism of protein secretion without clarifying the exact behavior and contribution of membrane phospholipids whose structural and metabolic dynamism can promote protein translocation through the cytoplasmic membrane.

For a more complete understanding of the mechanism of protein translocation through membranes, it is very important to find out whether the topogenic sequences of the preprotein interact with the phospholipids of the membranes. We were able to evaluate the contribution of translocation ATPase, the SecA protein, to protein-phospholipid interaction *in vivo*.

It was also possible to identify the features of the posttranslational modification of alkaline phosphatase in the absence of phosphatidylethanolamine in membranes and to determine the effect of PEA on the biosynthesis of alkaline phosphatase.

### References

- [1] M.V. Bogdanov, I.S. Kulaev, M.A. Nesmeyanova. «Studying a lipid - proteinaceous interactions in the process of transfer of proteins through a membrane of bacteria. Coordinate suppression by a protonoforms of synthesis and secretion exo-protein and exchange of phospholipids». *Biological membranes*. **1984**. Vol.1. P.495-502. (russian)
- [2] M.V. Bogdanov, N.E. Suzina, M.A. Nesmeyanova. «Particular qualities of phospholipid exchange and ultrastructural organization of cytoplasmic membrane *Escherichia coli* in the process of alkaline phosphatase secretion». *Biological membranes*. **1985**. Vol.2. P.367-375. (russian)
- [3] M.V. Bogdanov, G.N. Fedoseeva, M.A. Nesmeyanova. «Studying a lipid - proteinaceous interactions in the process of transfer of proteins through a membrane of bacteria. Exchange of phospholipids and content of periplasmic oligosaccharides in the process of biosynthesis and secretion of alkaline phosphatase in *Escherichia coli*». *Biological membranes*. **1986**. Vol.3. P.1241-1249. (russian)
- [4] O.A. Evdokimova, M.A. Nesmeyanova. «Phospholipid composition of cells and membranes *Escherichia coli* under repression and derepression of alkaline phosphatase biosynthesis». *Biochemistry*. **1977**. Vol.42. P.1791-1799. (russian)

- [5] O.A. Evdokimova, M.A. Nesmeyanova, I.S. Kulaev. «Induction of alkaline phosphatase synthesis in *Escherichia coli* by pre-incubation at low temperature». *Biochemistry*. **1978**. Vol.43. P.1680-1687. (russian)
- [6] O.A. Zemlyanukhina, V.V. Kolyecheva, M.A. Nesmeyanova. «Influence of lipotropic alcohols's agents on biosynthesis and repression of secreted alkaline phosphatase in *Escherichia coli*». *Biochemistry*. **1981**. Vol.46. P. 92-99. (russian)
- [7] S.N. Zolov, N.I. Mikhaleva, A.E. Kalinin, M.A. Nesmeyanova. «Interaction of prePhoA with phospholipids *in vivo* and *in vitro* depending on the charge of the N-terminus of the signal peptide and the content of anionic phospholipids in the membranes». *Biochemistry*. **2002**. Vol.67. P.1051-1060. (russian)
- [8] A.E. Kalinin, A.L. Karamyeshev, M.A. Nesmeyanova. «Violation of alkaline phosphatase processing, as a result of single amino acid substitutions, which effects on the composition and metabolism of phospholipids of *Escherichia coli* cells». *Biochemistry*. **1996**. Vol.61. P.104-113. (russian)
- [9] A.E. Kalinin, N.I. Mikhaleva, A.L. Karamyeshev, Z.N. Karamyesheva, M.A. Nesmeyanova. «Interaction of precursors of mutant alkaline phosphatases with membrane phospholipids *in vivo* and *in vitro*». *Biochemistry*. **1999**. Vol.64. P.1214-1223. (russian)
- [10] A.L. Karamyeshev, A.E. Kalinin, I.M. Tsfasman, V.N. Ksenzenko, M.A. Nesmeyanova. «Research of the biogenesis and secretion of alk aline phosphatase and its mutant forms in *Escherichia coli*. The effect of amino acid substitutions at the processing site and the *N*-terminal domain of the mature alkaline phosphatase polypeptide chain on its biogenesis». *Molecular biology*. **1994**. Vol.28. P.362-373. (russian)