

A new variant for the reduction of aromatic compounds until 2,5-dihydro derivatives by metal-alkylamine(ammonia)-aliphatic alcohol

© Pavel I. Fedorov,* Tatiana P. Fedorova, Oleg V. Ershov, Gennady P. Pavlov, Andrey N. Vasiliev, Vera A. Petrukhina, and Nikolay I. Koltsov[†]

Department of Organic and Pharmaceutical Chemistry. Department of Physical Chemistry and Macromolecular Compounds. Chuvash State University. I.N. Ulyanov. Moscow ave., 15.

Cheboksary, 428015. Chuvash Republic. Russia. Phone: +7 (352) 45-24-68. E-mail: paf@myrambler.ru

*Supervising author; [†]Corresponding author

Keywords: recovery, lithium alkylamine reduction by Benkeser, Birch recovery, reduction of phenol ethers.

Abstract

Development of methods for the preparation of aromatic hydrocarbons from aromatic compounds of di- and tetrahydro-derivatives, limiting and unsaturated alcohols and ketones of the mentane structure, further study of the reactions of the compounds obtained in the reactions of hydrohalogenation, halogenation, cationoid and anionoid transformations makes an important contribution to the chemistry of carbocyclic compounds and, first of all, to the chemistry of compounds series of Mentan. The cyclohexane compounds obtained from substituted benzene with two located at the 1,4-position, which differ markedly in reactivity by double bonds, can be used in the synthesis of hormones, prostaglandins, polyfunctional compounds with a wide range of biological effects. The compounds obtained can also be of interest for conformational analysis and dynamic stereochemistry of disubstituted cyclohexane. In connection with this, a new version of the reduction of aromatic compounds to 2,5-dihydro derivatives by metal-alkylamine (ammonia) – aliphatic alcohol systems is presented in this article. To restore various functional derivatives of benzene, a fundamentally new ratio of reagents in the reduction system (mole) is proposed: an aromatic compound : a metal : an aliphatic alcohol : an aliphatic amine (ammonia) 1 : 2-6 : 4-12 : 0.25-0.5. The recovery time of 1 mole of the initial arena is not more than 1 hour, the process temperature is 80-95 °C when converted to reaction products of 85-95%. The content of the reaction products of 2,5-dihydro derivatives is 50-80%, depending on the structure of the original arene. The proposed system is simpler in technological design than the known methods and lacks their main disadvantages: low temperature processes and use of no less than 20-30 moles of toxic, expensive amines per 1 mole arena or up to 200 moles of liquid ammonia (Birch method), or in its modification (the version of Wildes and Nelson) – addition of co-solvents.

References

- [1] A.J. Birch. The reduction of organic compounds by metal-ammonia solutions. *Quarterly Reviews*. **1950**. Vol.4. No.1. P.69-93.
- [2] E.A. Braude, A.A. Webb, M.U.S. Sultanawaba. Studies in the vitamin d field. Part III. Approaches to derivatives of 5-hydroxy-2-methylcyclohexanone. *J. Chem. Soc.* **1958**. P.3328-3336.
- [3] A.A. Akhrem, I.G. Reshetov, Yu.N. Titov. *Reactions and methods for studying organic compounds*. Edited by Kazansky, I.L. Knunyants, M.M. Shemyakin, N.N. Melnikov. *Moskow: Khimia*. **1969**. Vol.20. P.21-36. (russian)
- [4] R.A. Benkeser, J.A. Laugal, A. Rappa. *Process for reducing organic compounds with calcium, amine, and alcohol*. Patent USA 4533760(**1985**).
- [5] R.A. Benkeser, J.A. Laugal, B. Ganem. A safe and convenient new procedure for reducing aromatic compounds to birch-type products. *Tetrahedron Lett.* **1984**. Vol.25. No.20. P.2089-2092.
- [6] V.V. Bazylchik, P.I. Fedorov, T.N. Overchuk. *A method for producing piperitone*. Patent 658881. Publ. 02.28.**1980**. (russian)
- [7] A.L. Wilds, N.A. Nelson. The Facile Synthesis of 19-Nortestosterone and 19-Norandrostenedione from Estrone. *J. Am. Chem. Soc.* **1953**. Vol.75. No.21. P.5366-5369.
- [8] L. Reggel, R.A. Friedel, I. Wender. Lithium in Ethylenediamine: A New Reducing System for Organic

- Compounds. *J. Org. Chem.* **1957**. Vol.22. No.8. P.891-894.
- [9] V.V. Bazylchik, P.I. Fedorov, V.N. Odinkov. A new modification of the reduction of benzene and its alkyl substituted 1,4-dihydro derivatives in the lithium-isopropyl alcohol-ethylenediamine system. *Zhurn. organ. khim.* **1990**. Vol.26. P.1929-1932. (russian)
- [10] V.V. Bazylchik, T.N. Overchuk, P.I. Fedorov. Method for the preparation of 1-methoxycyclohex-1,4-diene or its methyl substituted. Patent 1685912. Publ. 10.23. **1991**. (russian)
- [11] P.I. Fedorov, T.P. Fedorova, V.P. Sheverdov, G.P. Pavlov, A.V. Eremkin. Investigation of compounds of the Mentan series. Synthesis of unsaturated primary alcohols of o- and p-mentane structure. *Zhurn. organ. khim.* **2016**. Vol.52. No.6. P.821-826. (russian)
- [12] R.A. Benkeser, M.A. Burrows, J.J. Hazdra, E.M. Kaiser. Reduction of Organic Compounds by Lithium in Low Molecular Weight Amines. VII. The Preparation of Dihydroaromatics. A Comparison of the Lithium-Amine and Birch Reduction Systems. *J. Org. Chem.* **1963**. Vol.28. No.4. P.1094-1097.
- [13] A.M. Ponomarev, A.N. Cherkasov, K.K. Pivnitsky. 1,2-Accumulation at reduction of aromatic compounds by alkali metals in ethers. *Zhurn. organ. khim.* **1971**. Vol.7. P.1428-1430. (russian)
- [14] P.I. Fedorov. *Synthesis, study of properties, chemical structure and transformations of o-mentadienes*. Author's abstract. dis. cand. chem. sciences: 02.00.03. Moscow. **1985**. 25p. (russian)
- [15] L. Fizer, M. Fizer. *Reagents for organic synthesis*. Moscow: Mir. **1970**. Vol.2. P.142. (russian)
- [16] A.J. Birch. Reduction by dissolving metals. Part VII. The reactivity of mesomeric anions in relation to the reduction of benzene rings. *J. Chem. Soc.* **1950**. P.1551-1556.
- [17] *The Aldrich library of infrared spectra*. By ed. Pouchert C.J. Milwaukee, Wisconsin. **1981**. 146p.
- [18] P.F. Schuda, S.J. Potlock, H. Ziffer. Use of the ketel claisen rearrangement for the synthesis of cyclic sesquiterpenes containing quaternary centers. A formal synthesis of cuparene. *Tetrahedron*. **1987**. Vol.43. No.3. P.463-468.
- [19] E. Taskinen, K. Nummelin. Thermodynamics of vinyl ethers. 31. Isomer equilibria in some six- and seven-membered cyclic dienes. *J. Org. Chem.* **1985**. Vol.50. No.24. P.4844-4847.
- [20] G.L. Burge, P.J. Collins, J.D. Reitze. Synthesis of 3-Ethenyl-4-methylcyclohex-2-en-1-one from 4-methylanisole. *Austr. J. Chem.* **1982**. Vol. 35. No.9. P.1913-1925.