

Immobilization of pyridinecarboxylic acids in a polymer nanocontainer based on the AB-17-8 strongly basic anion exchanger

© Heinrich N. Altshuler,* Galina Yu. Shkurenko,[†]
Natalia V. Malyschenko, and Sergey Yu. Lyrschikov

Institute of Coal Chemistry and Material Science. Federal Research Center of Coal and Coal Chemistry.

Siberian Branch of Russian Academy of Sciences. Sovetsky Ave., 18. Kemerovo, 650000. Russia.

Phone: +7 (384-2) 36-88-04. E-mail: altshulerh@gmail.com

*Supervising author; [†]Corresponding author

Keywords: pyridinecarboxylic acids, AB-17-8 anionite, immobilization.

Abstract

The possibility of the immobilization of pyridinecarboxylic (nicotinic, isonicotinic, dipicolinic, and cinchomeronic) acids in the phase of AB-17-8 strongly basic anion exchanger and subsequent their release from the nanocontainer into the solution has been researched. The immobilization of pyridinecarboxylic acids in the AB-17-8 polymer nanocontainer was performed by ion exchange sorption. In the ¹³C NMR spectra of solid-state samples of the AB-17-8 anionite in the OH, in dipicolinic and cinchomeronic acids are observed the resonance absorption bands of ¹³C in the ranges of chem. shifts of 40-70, 120-150 and 190-210 ppm, corresponding to the carbon of the elementary unit of the anion exchanger. In the NMR spectrum of the anion exchanger containing the immobilized dipicolinic and cinchomeronic acids, an additional resonance line (170 ppm) appears that corresponds to the carbon of two carboxyl groups of pyridinecarboxylic acids. This is absent in the anionite spectrum in the OH form. From the analysis of NMR spectra it follows that pyridinecarboxylic acids were encapsulated in a nanocontainer based on the AB-17-8 anionite. The dynamic ion-exchange capacities of strongly basic anionite during the sorption (immobilization) of pyridinecarboxylic acids in a polymer nanocontainer coincides with the ion-exchange capacities of the anionite after desorption (release) of these acids from the polymer nanocontainer. The ion-exchange capacity of a polymer nanocontainer based on the anionite for nicotinic acid is 2.8 meq/g, and for isonicotinic acid is 2.7 meq/g. This corresponds to the content of ionogenic groups in the polymer. The ion exchange capacities of the anionite for dipicolinic acid and cinchomeronic acid are 2.0 meq/g and 2.1 meq/g, respectively. It is shown that the kinetics of the processes of the immobilization of pyridinecarboxylic acids in a polymer nanocontainer based on the AB-17-8 strongly basic anion exchanger and their release is determined by the diffusion of the organic components in the polymeric phase.

References

- [1] D.J. Cram, S.K. Karbach, Kim Hye-Eyn, C.B. Knoler, E.F. Maverick, J.L. Ericson, R.C. Helgeson. Host-guest complexation. 46. Cavitands as open molecular vassels form solvates. *J. Am. Chem. Soc.* **1988**. Vol.110. P. 2229-2237.
- [2] H.N. Al'tshuler, Yu.A. Makarov, Yu.S. Porodnova. Adsorption of novocaine and anesthesin on KU-2 and KB-4P-2 cationites. *Pharmaceutical Chemistry Journal.* **1969**. Vol.3. No.7. P.404-406. (russian)
- [3] H.N. Altshuler, G.Yu. Shkurenko, S.Yu. Lyrschikov, A.A. Gorlov, and O.H. Altshuler. Solid nanoreactor. Part 5. Polymer nanocontainers for benzocaine. *Butlerov Communications.* **2015**. Vol.44. No.11. P.69-72. ROI: jbc-02/15-44-11-69
- [4] E.V. Ostapova, G.Yu. Shkurenko, S.Yu. Lyrschikov, and H.N. Altshuler. Sulphonated network polymers as containers for bioactive substances. *Butlerov Communications.* **2016**. Vol.48. No.10. P.37-42. ROI: jbc-01/16-48-10-37
- [5] *The International Pharmacopoeia*, Fifth Edition (incl. First, Second, Third and Forth Supplements). Available at: [http://apps.who.int/phint/en/p/docf/].
- [6] M.D. Mashkovcky. *Pharmaceutical. Moscow: New wave.* **2002**. Vol.1. Iss.14. 608p. (russian)
- [7] Black Gordon, Depp Elizabeth, and Corson B. B. Oxidation of certain methylpyridines to pyridine carboxylic acids. *J. Org. Chem.* **1949**. Vol.14. No.1. P.14-21.

- [8] V.V. Veselovsky, L.L. Danilov, S.D. Maltsev, A.V. Pronin, A.N. Narovlyansky, A.V. Sanin, A.V. Deeva, A.M. Amchenkova. 1: 2 complex of germanium and 2,6-pyridinedicarboxylic acid, a process for its preparation, a pharmaceutical composition. *RU2171259C2*. 27.07. **2001**. <http://ru-patent.info/21/70-74/2171259.html>
- [9] A.F. Lindenstruth, C.A. VanderWerf. The Preparation of Quinolinic and Cinchomeric Acids by Ozone Oxidation. *J. of Amer. Chem.Soc.* **1949**. Vol.71. No.9. P.3020-3021.
- [10] O.N. Phedoseeva, E.P. Cherneva, N.N. Tunitskii. Investigation of the kinetics of ion exchange sorption. *Russian Journal of Physical Chemistry*. **1959**. Vol.33. No.4. P.936-938. (russian)
- [11] L.K. Shataeva, N.N. Kuznetsova, G.E. Elkin. Carboxyl ion exchangers in biology. *Leningrad: Science*. **1979**. 286p. (russian)