

## Synthesis of alkylphenols lactamomethyl derivatives

© Stepan V. Vorobyev,<sup>+</sup> Olga V. Primerova, Vladimir N. Koshelev,\* and Lyudmila V. Ivanova

Department of Organic Chemistry & Petroleum Chemistry, Gubkin Russian State University of Oil and Gas (National Research University). Leninsky Prospekt, 65. Moscow, 119991. Russia. Phone: +7 (916) 641-72-16.

E-mail: vorstepan@yandex.ru

\*Supervising author, <sup>+</sup>Corresponding author

**Keywords:** alkylphenols, organic synthesis, QSAR, quantum-chemical calculations.

### Abstract

In this paper we describe the synthesis of lactamomethyl derivatives of several alkylphenols (thymol and 2,4-di-*t*-butyl-phenol) with butyrolactam, valerolactam, caprolactam and 4-phenylpyrrolidone fragments. The development of such compounds is one of the modern trends for the search of new and effective antioxidants with low toxicity and, as a result, with wide range of possible usage. The structures of target compounds were confirmed by IR and NMR study. In IR spectra there are carbonyl group peaks in lower frequencies (about 1600 cm<sup>-1</sup>) than expected due to the formation of inter- and intramolecular hydrogen bonds between this group and phenolic hydroxyl group. In <sup>1</sup>H NMR spectra the signals of protons of lactam fragments and phenolic alkyl substituents are located in the high field region (1.50-3.50 ppm) and have a form of corresponding multiplets. The value of the chemical shift of the "methylene bridge" (-CH<sub>2</sub>-group) protons is about 4.00 ppm. This signal is a singlet in most cases. The signals of aromatic and hydroxylic protons are located in the weak-field region. For several products, containing a fragment of 4-phenylpyrrolidone, the signal of the methylene bridge is an AB system due to the presence of asymmetric nitrogen atom. *In silico* estimation of possible pharmacological effects and acute rat toxicity for target compounds was carried out using PASS-Online and GUSAR-Online services. It was shown, that studied compounds are expected to be nontoxic or low toxic (Class 5 or 4 of rat acute toxicity) and possess wide range of possible biological effects and. Energy of dissociation of ArO-H bond was calculated to reveal possible antioxidant activity of target compounds using quantum chemical method (semi-empirical PM6) in ROO· + ArOH → ROOH + ArO· reaction model. According to the calculation results, introducing of lactamomethyl fragment in the *para*-position of phenolic ring decreases energy of the dissociation of ArO-H bond. In case of *ortho*-substitution, increasing of the energy was observed due to the formation of hydrogen bond between phenolic hydroxyl group and lactam carbonyl group.

### References

- [1] G.W. Burton, K.U. Ingold. Autoxidation of biological molecules. 1. Antioxidant activity of vitamin E and related chain-breaking phenolic antioxidants in vitro. *J. Am. Chem. Soc.* **1981**. Vol.103. No.21. P.6472-6477. DOI:10.1021/ja00411a035
- [2] F. Iverson. Phenolic antioxidants: Health Protection Branch studies on butylated hydroxyanisole. *Cancer letters.* **1995**. Vol.93. No.1. P.49-54. DOI:10.1016/0304-3835(95)03787-W
- [3] K.U. Ingold. Inhibition of the autoxidation of organic substances in the liquid phase. *Chem. Rev.* **1961**. Vol.61. P.563-589. DOI: 10.1021/cr60214a002
- [4] S.K. Filoche, K. Soma, C.H. Sissons. Antimicrobial effects of essential oils in combination with chlorhexidine digluconate. *Oral Microbiol. Immunol.* **2005**. Vol.20(4): 221-225. doi:10.1111/j.1399-302X.2005.00216.x
- [5] D. Hu, J. Coats. Evaluation of the environmental fate of thymol and phenethyl propionate in the laboratory. *Pest Manag. Sci.* **2008**. Vol.64(7): 775-779. doi:10.1002/ps.1555
- [6] V.N. Koshelev, V.I. Kelarev, N.V. Belov, O.V. Malova, S.L. Osipov, V.G. Spirkin. Effect of azoles and sym-triazines with hindered phenol fragments on protective properties of turbine oils. *Chemistry and technology of fuels and oils.* **1995**. Vol.31. No.1. P.26-29.
- [7] V.N. Koshelev, I.A. Golubeva, E.V. Klinaeva, V.I. Kelarev. Stabilization of ecologically clean diesel fuel by means of combinations of additives. *Chemistry and technology of fuels and oils.* **1996**. Vol.32. No.4. P.189-194.

- [8] V.I. Kelarev, M.A. Silin, I.A. Golubeva, O.A. Borisova. Stabilization of distillate fuels in storage. *Chemistry and technology of fuels and oils*. **2000**. Vol.36. No.2. P.111-115.
- [9] K. Katayama, H. Shiota, S. Kobayashi, K. Terato, H. Ikuta, I. Yamatsu. *In vitro* effect of N-methoxy-3-(3, 5-ditert-butyl-4-hydroxybenzylidene)-2-pyrrolidone (E-5110), a novel nonsteroidal anti-inflammatory agent, on generation of some inflammatory mediators. *Agents and actions*. **1987**. Vol.21. No.3-4. P.269.
- [10] T. Hidaka, K. Hosoe, T. Yamashita, K. Watanabe, Y. Hiramatsu, H. Fujimura. Analgesic and anti-inflammatory activities in rats of  $\alpha$ -(3,5-di-t-butyl-4-hydroxybenzylidene)- $\gamma$ -butyrolactone (KME-4), and its intestinal damage. *Journal of pharmacy and pharmacology*. **1986**. Vol.38. No.10. P.748-753.
- [11] S.V. Satheesh, A.V. Radha, K.K.N. Giriya, K.N. Rajasekharan, P.R. Maheswari. Hindered phenolic aminothiazoles – Synthesis,  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibitory and antioxidant activities. *Journal of the Serbian Chemical Society*. **2017**. Vol.82. No.10. P.1087.
- [12] A.H. Gouliaev, A. Senning. Piracetam and other structurally related nootropics. *Brain Research Reviews*. **1994**. Vol.19. No.2. P.180-222. DOI: 10.1016/0165-0173(94)90011-6.
- [13] H. Mohrle, C. Miller. Lactambildung bei Amindehydrierungen ohne Nachbargruppenbeteiligung. *Arch. Pharm. (Weinheim)*. **1983**. Vol.316. P.160.
- [14] DE 2616374
- [15] WO2005054188A1
- [16] US3829528
- [17] US3136766
- [18] DE 134979
- [19] A. Einhorn, T. Mauermayer, C. Ladisch, G. Schupp. Über die N-Methylolverbindungen der Säureamide. *Annalen*. **1905**. P.207-305.
- [20] J. Barry, E. Mayeda, S. Ross. The amidoalkylation of aromatic hydrocarbons. *Tetrahedron*. **1976**. Vol.33. P.369-372.
- [21] A. Sadym, A. Lagunin, D. Filimonov, V. Poroikov. *SAR QSAR Environ. Res*. **2003**. Vol.14. P.339.
- [22] D.A. Filimonov, A.A. Lagunin, T.A. Glorizova, A.V. Rudik, D.S. Druzhilovski, P.V. Pogodin, V.V. Poroikov. *Chem. Heterocycl. Compd*. **2014**. Vol.50(3). P.444.
- [23] A. Lagunin, A. Zakharov, D. Filimonov, V. Poroikov. *Mol. Inf*. **2011**. Vol.30(2-3). P.241.
- [24] Васильев Р.Ф., Кънчева В.Д., Федорова Г.Ф., Бътовска Д.И., Трофимов А.В. Антиоксидантная активность халконов. Хемилюминесцентное определение реакционной способности и квантово-химический расчет энергий и строения реагентов и интермедиатов. *Кинетика и катализ*. **2010**. Т.51. №4. С.533-541.
- [25] Gaussian 09, Revision **D.01**, M.J. Frisch; G.W. Trucks; H.B. Schlegel; G.E. Scuseria; M.A. Robb; J.R. Cheeseman; G. Scalmani; V. Barone; B. Mennucci; G.A. Petersson; H. Nakatsuji; M. Caricato; X. Li; H.P. Hratchian; A.F. Izmaylov; J. Bloino; G. Zheng; J.L. Sonnenberg; M. Hada; M. Ehara; K. Toyota; R. Fukuda; J. Hasegawa; M. Ishida; T. Nakajima; Y. Honda; O. Kitao; H. Nakai; T. Vreven; J.A. Montgomery, Jr.; J.E. Peralta; F. Ogliaro; M. Bearpark; J.J. Heyd; E. Brothers; K.N. Kudin; V.N. Staroverov; R. Kobayashi; J. Normand; K. Raghavachari; A. Rendell; J.C. Burant; S.S. Iyengar; J. Tomasi; M. Cossi; N. Rega; J.M. Millam; M. Klene; J.E. Knox; J.B. Cross; V. Bakken; C. Adamo; J. Jaramillo; R. Gomperts; R.E. Stratmann; O. Yazyev; A.J. Austin; R. Cammi; C. Pomelli; J.W. Ochterski; R.L. Martin; K. Morokuma; V.G. Zakrzewski; G.A. Voth; P. Salvador; J.J. Dannenberg; S. Dapprich; A.D. Daniels; Ö. Farkas; J.B. Foresman; J.V. Ortiz; J. Cioslowski; D.J. Fox. *Gaussian, Inc., Wallingford CT*. **2009**.