

Synthesis of benzyloxybenzyl derivatives of 2- and 4-(1*H*-azol-1-ylmethyl)phenols

© Zoya P. Belousova,¹⁺ Pyotr P. Purygin,¹ and Vladimir N. Koshelev²

¹ Department of Organic, Bioorganic and Medicinal Chemistry. Samara National Research University. Moskovskoe shosse, 34. Samara, 443086. Samara Region. Russia.

Phone: +7 (846) 334-54-59. E-mail: zbelousova@mail.ru

² Department of Organic Chemistry and Chemistry of Oil. Gubkin Russian State University of Oil and Gas (National Research University). Leninsky Ave., 65, korp. 1. Moscow, 119991. Russia.

*Supervising author; ⁺Corresponding author

Keywords: heterocycles, imidazole, 2-methylimidazole, 1,2,4-triazole, benzimidazole, 2-methylbenzimidazole, benzotriazole, 2-(1*H*-azol-1-ylmethyl)phenols, 4-(1*H*-azol-1-ylmethyl)phenols, benzyl phenyl ethers of 1*H*-azoles, 1-[2-(benzyloxy)benzyl]-1*H*-azoles and 1-[4-(benzyloxy)benzyl]-1*H*-azoles, lipophilicity.

Abstract

The synthesis of 1-[2-(benzyloxy)benzyl]- and 1-[4-(benzyloxy)benzyl]-1*H*-azoles has been reported. These compounds were prepared from the corresponding 2- and 4-(1*H*-azol-1-ylmethyl) phenols in the presence of sodium hydride in anhydrous dimethylformamide. As the nitrogen component, nitrogen compounds with different nitrogen atoms in the cycle, their methylated and benzo-fused derivatives: imidazole, 2-methylimidazole, 1,2,4-triazole, benzimidazole, 2-methylbenzimidazole, benzotriazole were used. The structure of the compounds was confirmed by IR and ¹H NMR spectroscopy. The band of stretching vibrations of the hydroxyl group in the region 3135-2480 cm⁻¹ in the IR spectra of 1-[2-(benzyloxy)benzyl]- and 1-[4-(benzyloxy)benzyl]-1*H*-azoles characteristic of azolylmethylphenols, the link is recorded in the form of several peaks in the 1260-1014 cm⁻¹ region, which proves the presence of 2- and 4-(1*H*-azol-1-ylmethyl)phenols in the structure of the benzyloxybenzyl derivatives of the fragment (–C_{apom}–O–C–). There are additional signals in the form of singlets in the range 5.12 ppm for *ortho*-derivatives and 5.14 ppm for *para*-derivatives, respectively, for 1-[2-(benzyloxy)benzyl]- and 1-[4-(benzyloxy)benzyl]-1*H*-azoles in the ¹H NMR spectra for the oxymethylene groups –CH₂O–. The ALOGPS 2.1 program was used to calculate the lipophilicity of both the obtained and the original compounds. All 1-[2-(benzyloxy)benzyl]- and 1-[4-(benzyloxy)benzyl]-1*H*-azoles, in contrast to 2- and 4-(1*H*-azol-1-ylmethyl) phenols are characterized by high lipophilicity. The lipophilicity of benzyl esters of 4-(1*H*-azol-1-ylmethyl) phenols is higher than that of 2-(1*H*-azol-1-ylmethyl)phenols, which is probably due to the greater spatial availability of the hydrophobic van der Waals surface in benzyl ester of 4-(1*H*-azol-1-ylmethyl)phenols. Methylated azole fragments, derivatives are also characterized by increased lipophilicity in comparison with their unmethylated analogues. The benzimidazole and benzotriazole derivatives were most lipophilic.

References

- [1] M.D. Mashkovskii. Drugs. Moscow: Novaya Volna. 2012. 931p. (russian)
- [2] Justin I. Montgomery, Peter L. Toolgool, Kim M. Hutching, Jia Liu, Lakshmi Narasimhan, Timothy Braden, Michael R. Dermeyer, Angela D. Kulynych, Yvonne D. Smith, Joseph S. Warmus, Clarke Taylor. Discovery and SAR of benzyl phenyl ester as inhibitors of bacterial phenylalanyl-tRNA synthetase. *Bioorganic & Medical Chemical Letters*. 2009. Vol.19. P.665-669.
- [3] John Lawrence Andreassi, II, Wilmington, Andrew Edmund. Heterobicycle-substituted azolyl benzene fungicides. *Patent US 8,921,360 B2*. 2014.
- [4] Massimo Pinori, Paderno d'Adda Lecco, Maria Lattanzio, Daniela Modena, Paolo Mascagni. Azole derivative useful as antifungal agents with reduced interaction with metabolic cytochromes. *Patent US 7,427,620 B2*. 2008.
- [5] O.A. Kolyamshin, Yu.N. Mitrasov, V.A. Danilov, I.N. Smolina, and N.I. Koltsov. Synthesis of gem-dichlorocyclopropylmethylthio-substituted derivative benzimidazoles, benzoxazoles, 4,6-

- SYNTHESIS OF BENZYLOXYBENZYL DERIVATIVES OF 2- H 4-(1H-AZOL-1-YLMETHYL)PHENOLS* _____ 132-137
dimethylpyrimidines and 1,2,4-triazoles. *Butlerov Communications*. **2016**. Vol.45. No.3. P.6-10. ROI:
jbc-02/16-45-3-6
- [6] A. Safronova, E.A. Teplova, R.V. Shafigulin, Z.P. Belousova, A.V. Bulanova. Thermodynamics of Sorption of Some Benzimidazole Derivatives on a Nonpolar Sorbent from Aqueous-Acetonitrile Solution *Fizikokhimiya Poverkhnosti i Zashchita Materialov*. **2014**. Vol.50. No.1. P.38-41.
- [7] I.N. Karaseva, E.I. Demchenko, and S.V. Kurbatova. 1,2,4-Triazine and 1,2,4-triazole derivatives sorption under condition of high performance liquid chromatography. *Butlerov Communications*. **2018**. Vol.54. No.5. P.40-49. ROI: jbc-02/18-54-5-40
- [8] V.A. Osyanin, P.P. Purygin, Z.P. Belousova, P.E. Krasnikov. Synthesis and glycosilation of 4-(1H-azol-1-yl-methyl)phenols. *Proceedings of high schools. Ser. Chemistry and Chemical Technology*. **2003**. Vol.46. Iss.1. P.138-141. (russian)
- [9] V.A. Osyanin, P.P. Purygin, Z.P. Belousova. The interaction of salicylic alcohol with certain azoles. *Proceedings of high schools. Ser. Chemistry and Chemical Technology*. **2003**. Vol.46. Iss.6. P.23-24. (russian)
- [10] <http://www.vcclab.org/lab/alogps/>