

To the question of standardization of the polyextract dry “*Hippophae-5*” having adaptogeny and anti-inflammatory activity

© Tatiana V. Kornopoltseva,¹ Elena A. Botoeva,² Evgeny V. Petrov,¹ and Julia Shurigina³

¹Laboratory of Biomedical Research. Institute of General and Experimental Biology.

Sakhyanovoy St., 6. Ulan-Ude. Russia. Phone: +7 (9021) 64-55-81. E-mail: tv-kornopol@mail.ru

²Department of Obstetrics and Gynecology with the Course of Pediatrics. Buryat State University. Smolin St., 24a. Ulan-Ude. Russia.

³Department Social Technologies of the East Siberian State University of Technologies and Management. Klyuchevskaya St., 40 century of Ulan-Ude. Russia.

*Supervising author; †Corresponding author

Keywords: biologically active substances, standardization, polyextract.

Abstract

In an arsenal of the Tibetan medicine at diseases of respiratory organs multicomponent medicines which part plants from flora of *Byryatia* are used. Collecting (the conditional name “*Hippophae-5*”) includes the following herbs: fruits of *Hippophaerhamnoides* L.) fruits of *Piper longum* L. root *Glycyrrhizauralensis* Fisch., root *Phlojodicarpussibiricus* (*Stephan ex Spreng*) Koso-Pol. Tausch (roots), fruits *Malussylvestris* Mill. There are components of anti-inflammatory action a *Glycyrrhiza*, *MalusPhlojodicarpus*; antioxidant – a *Glycyrrhiza*, *Hippophae*, *Piper*, *Malus*; immunomodulatory – a *Glycyrrhiza*, *Malus*; antibacterial – the *Glycyrrhiza*, a *Hippophae* promoting increase in resistance of an organism. The rematseration method taking into account optimum parameters of extraction has received the complex polyextract consisting of dry extracts of fruits of a *Hippophae hamnoides* L., fruits of *Pepper*, roots of a *Glycyrrhizal*, roots of a *Phlojodicarpus* and fruits of *Malus* the mass parts 2:2:1:1:1 taken in the ratio, respectively. The HLH method in polyextract dry has installed existence of six marker components which raw source is the *Phlojodicarpus* (visnadin), *Pepper* (piperin), a *Glycyrrhiza* (likviritin, glycyrrhizic acid), *Malus* (catechin). The dominating compounds of *Hippophae-5* extract, are glycyrrhizic acid which content was 27.4 mg/g; on a share of benzopyrones (visnadin), 7.65 mg/g from the mass of dry extract, are the share of a share of a piperin – 2.23 mg/g, respectively and of a catechin share – 0.52 mg/g. The technique of quantitative determination of content of glycyrrhizic acid is developed (24.15%) in polyextract dry which can be used for standardization of this object.

Литература

- [1] State pharmacopeia of the USSR. XI prod. Moscow. 1990. Iss.2. P.364-365. (russian)
- [2] M.V. Egorov, V.A. Kurkin, G.G. Zapesochnaya, V.A. Bikov. Validation of techniques of the qualitative analysis of raw materials and medicines of a glycyrrhiza. *Pharmacy*. 2005. Vol.53. No.1. P.9-12. (russian)
- [3] M.V. Egorov, V.A. Kurkin. Improvement of methods of standardization of roots of a glycyrrhiza. *News of the Samara scientific center of the Russian Academy of Sciences*. 2011. Vol.13. No.1(8). P.1992-1995. (russian)
- [4] Vegetable resources of Russia: Wild-growing flowering plants, their component structure and biological activity. In 6 t. Vol.2. Families Actinidiaceae-Malvaceae, Euphorbiaceae-Haloragacea. autor L.M. Belenovskaya, E.E. Lesiovskaya, N.S. Bobyliova. *St. Petersburg. –Moscow*. 2009. 513p. (russian)
- [5] Vegetable resources of Russia: Wild-growing flowering plants, their component structure and biological activity. In 6 t. Vol.3. Fabaceae-Apiaceae.sost families. L.M. Belenovskaya, E.E. Lesiovskaya, N.S. Bobyliova. *St. Petersburg. –Moscow*. 2010. 601p. (russian)
- [6] Vegetable resources of Russia: Wild-growing flowering plants, their component structure and biological activity. In 6 t. Vol. 5. 2. Asteraceae (Compositae). Part 1. Rodyachillea-Youngia. autor Belenovskaya L.M., Lesiovskaya E.E., Bobylevan. *St. Petersburg. –Moscow*. 2013. 312p. (russian)
- [7] Sanchzhay-chzhamtso. Vaydurya-onbo: comment of Chzhud-shi. Desridsanchzhay-chzhamtso; the lane with Tibetan. Note D.B. Dashiyevev's articles. *Moscow: Eastern Literature*. 2014. P.541. (Library of the Tibetan medicine). (russian)

- [8] Sumati Pradzhnya. Kunpan-Dudz (Amrita extract, useful to all). Big prescription reference book of the Agin datsan. Sumatipradzhnya. Per about Tibet., note, decree. D.B. Dashiyeva. *Moscow: Eastern Literature*. **2008**. 214p. (russian)
- [9] Chzhud-sha: canon of the Tibetan medicine the lane with Tib., note., D.B. Dashiyev's indexes. *Moscow: Westlit-RAS Publishing House*. **2001**. P.230. (russian)
- [10] R.K. Choudhary, P.L. Swarnkar. Antioxidant activity of phenolic and flavonoid compounds in some medicinal plants of India. *Natural Product Research*. **2011**. Vol.25. No.11. P.1101-1109.
- [11] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova. A Novel HPLC-Assisted Method for Investigation of the Fe²⁺-Chelating Activity of Flavonoids and Plant Extracts – 2014. *Molecules*. **2014**. Vol.19. P.18296-18316.
- [12] D.N. Olennikov, N.I. Kashchenko, N.K. Chirikova, S.S. Kuz'mina. Phenolic Profile of *Potentilla anserina* L. (Rosaceae) Herb of Siberian Origin and Development of a Rapid Method for Simultaneous Determination of Major Phenolics in *P. anserina* Pharmaceutical Products by Microcolumn RP-HPLC-UV – 2015. *Molecules*. **2015**. Vol.20. P.224-248.
- [13] J. Han, G. Borjihan, R. Bai, Z. Sun, N. Bao, X. Chen, X. Jing. Synthesis and anti-hyperlipidemic activity of a novel starch piperinic ester. *Carbohydrate Polymers*. **2008**. Vol.71. No.3. P.441-447.