

## Reaction of 4,4'-bis(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-3,3'-dimethoxydiphenylmethane with 2-furylmethanol

© Yury N. Mitrasov,<sup>1\*</sup> Svetlana P. Yashkildina,<sup>2</sup> Oksana V. Kondratieva,<sup>1</sup>  
Oleg A. Kolyamshin,<sup>3</sup> and Vadim A. Kozlov<sup>3</sup>

<sup>1</sup> Department of Bioecology and chemistry. I. Yakovlev Chuvash State Pedagogical University.  
K. Marx St., 38. Cheboksary, 428000. Chuvash Republic. Russia.

Phone: +7 (8352) 22-02-28. E-mail: mitrasov\_un@mail.ru.

<sup>2</sup> Department of Humanities, Natural Sciences and General Professional Disciplines. Branch of Ukhta State Technical University. Neftyanikov St., 33. Usinsk, 169710. Republic of Komi. Russia.

Phone: +7 (82144) 29-1-74. E-mail: svetlana-jashkildina@rambler.ru

<sup>3</sup> Department of Medical Biology with Course of Microbiology and Virology. I.N. Ulyanov Chuvash State University. Moskovsky Ave., 15. Cheboksary, 428015. Chuvash Republic. Russia.

Phone: +7 (8352) 45-26-97. E-mail: pooh12@yandex.ru

\*Supervising author; †Corresponding author

**Keywords:** 4,4'-methylene-bis-(*o*-anisidine), maleic anhydride, 4,4'-bis(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-3,3'-dimethoxydiphenylmethane, 2-furylmethanol, diene synthesis, fluorescence.

### Abstract

The reaction of 4,4'-bis(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-3,3'-dimethoxydiphenylmethane with 2-furylmethanol is described. The relevance of the study is due to the fact that 2,5-dihydropyrrole-2,5-diones (maleic acid imides), due to the presence of a highly active double bond, readily enter polymerization and copolymerization reactions with various unsaturated monomers, various nucleophilic and electrophilic reagents are attached, participate in a number of cycloaddition reactions, as well as substitution. Along with this many maleimide derivatives have biological activity. Thus they are widely used in agriculture as pesticides, because they exhibit high insecticidal, fungicidal and herbicidal activities. Maleimides are also proposed as highly effective pharmaceutical preparations for the treatment of a number of diseases, including cardiovascular diseases, Alzheimer's disease, type 2 diabetes, cancer and HIV. The starting 4,4'-bis(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-3,3'-dimethoxydiphenylmethane was synthesized by a two-step methodic method based on 4,4'-methylene-bis-(*o*-anisidine). Firstly, 4,4'-bis[*N*-(2-carboxyethenyl)carbomoyl]-3,3'-dimethoxydiphenyl-methane was prepared by reacting diamine with maleic anhydride in a molar ratio of 2: 1 in acetone, which under heating with acetic anhydride in the presence of anhydrous sodium acetate undergoes intramolecular cleavage of water with the formation of diimide. Reactions of 4,4'-bis(2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)-3,3'-dimethoxydiphenylmethane with 2-furylmethanol were carried out at a molar ratio of reagents 1 : 1 and 1:2 at room temperature. Absolute 1,4-dioxane was used as the solvent. The progress of the reactions was monitored by thin layer chromatography. IR and <sup>1</sup>H NMR spectroscopy showed that the reactions proceed according to the scheme of [4+2]-cycloaddition and lead to the formation of mono- or bis-adducts of diene synthesis. The phenomenon of induced fluorescence for synthesized adducts was revealed.

### References

- [1] O.A. Kolyamshin, V.A. Danilov, S.Yu. Vasilieva. Maleinimide: synthesis, properties and biological activity. *Dep. VINITI*, No. 250-B2010 from 06.05.2010. 75p. (russian)
- [2] Yu.N. Mitrasov, A.A. Avruyskaya, L.M. Sadikova, O.V. Kondratyeva, O.A. Kolyamshin. The reaction of *N*-aryl-2,5-dihydropyrrol-2,5-diones with functional derivatives of furan. *Cheboksary: Chuvash. State Pedagogical University*. 2013. 155p. (russian)
- [3] Yu.N. Mitrasov, O.A. Kolyamshin, V.A. Danilov. Maleimides: synthesis, properties and polymers on their basis. *Cheboksary: Chuvash. State Pedagogical University*. 2017. 286p. (russian)
- [4] O.A. Kolyamshin, Yu.N. Mitrasov, V.A. Danilov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted nitrogen-containing heterocycles. Part 1. Synthesis of *gem*-dichlorocyclopropyl-methylthiosubstituted derivative benzimidazoles, benzoxazoles, 4,6-dimethylpyrimidines and 1,2,4-triazoles. *Butlerov Communications*. 2016. Vol.45. No.3. P.6-10. DOI: 10.37952/ROI-jbc-01/16-45-3-6

- REACTION OF 4,4'-BIS (2,5-DIHYDRO-2,5-DIOXO-1H-PYRROL-1-YL)-3,3'-DIMETHOXYDIPHENYLMETHANE... \_\_ 50-55
- [5] O.A. Kolyamshin, Yu.N. Mitrasov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted nitrogen-containing heterocycles. Part 2. Synthesis and properties of 1-(2,2-dichlorocyclopropylmethyl)pyrazoles and 5-pyrazolones. *Butlerov Communications*. **2016**. Vol.46. No.4. P.77-80. DOI: 10.37952/ROI-jbc-01/16-46-4-77
- [6] O.A. Kolyamshin, Yu.N. Mitrasov, I.N. Smolina, N.I. Koltsov. *gem*-Dichlorocyclopropane substituted nitrogen-containing heterocycles. Part 3. Synthesis and properties of 3-[(1-methyl-2,2-dichlorocyclopropyl)methyl]hydantoin. *Butlerov Communications*. **2016**. Vol.48. No.10. P.103-107. DOI: 10.37952/ROI-jbc-01/16-48-10-103
- [7] A.c. 390078 (USSR), IPC C 07 C 101/20. Method for the preparation of aminodicarboxylic acids. I.O. Elin, A.A. Artyushina, Yu.Ya. Meshcheryakov, V.P. Menshutin, T.M. Abramova, M.S. Akutin. – Declared. 23.03.71; Publ. 29.11.73. *AJChem*. **1974**. 15N181P (russian)
- [8] V.A. Kozlov, Yu.N. Mitrasov, O.V. Kondrat'eva, I.V., Gordeeva, O.B. Polyakova, O.E. Kirillova, A.N. Kochneva, A.A. Avruyskaya, V.A. Sorokina, O.A. Kolyamshin. Photometry 4-Aza-4-aryl-1-hidroksimethyl-10-oxa-3,5-dioxotricyclo[5.3.1<sup>1.7</sup>.0<sup>2.6</sup>]dec-8-enes. *Materials of the seventh international scientific school "Science and innovations – 2012" ISS "SI-2012". Yoshkar-Ola: Mari State University*. **2012**. P.232-235. (russian)
- [9] Yu.N. Mitrasov, A.A. Avruyskaya, O.V. Kondratieva. The interaction of the nitro- and aminosubstituted of phenylmaleimide and furfurylated alcohol. *Journal of General Chemistry*. **2015**. Vol.85. No.1. P.82-85. (russian)
- [10] Yu.N. Mitrasov, A.A. Avruyskaya, O.B. Polyakova, O.E. Ivanova. Interaction derivative (2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl)benzoic acids  $\alpha$ -furylmethanol. *Journal of Organic Chemistry*. **2015**. Vol.51. No.8. P.1206-1207. (russian)
- [11] Yu.N. Mitrasov, I.V. Gordeyeva, O.V. Kondratyeva, A.A. Avruyskaya, O.E. Kirillova. Reactions of derivatives of 2,5-dihydro-2,5-dioxo-1H-pyrrol-1-ylarenecarboxylic(-phosphonic) acids and furfuryl alcohol. *Bulletin of the Chuvash State Pedagogical University of I.Ya. Yakovlev*. **2012**. No.2 (74). Part1. P.104-107. (russian)
- [12] V.A. Kozlov, Yu.N. Mitrasov, A.A. Avruyskaya, O.B. Polyakova, O.V. Kondratieva. Acute toxicity of derivatives of *m* - and *p*-(4-Aza-1-hidroksimethyl-10-oxa-3,5-dioxotricyclo[5.2.1<sup>1.7</sup>.0<sup>2.6</sup>]dec-8-en-4-yl)-benzoic acids. *The ninth international scientific school "Science and innovations – 2014" ISS "SI-2014". Yoshkar-Ola: Mari State University*. **2014**. P.183-185. (russian)
- [13] Yu.N. Mitrasov, A.A. Avruyskaya, O.V. Kondratieva, V.A. Kozlov. The reaction of *N*-aryl-2,5-dihydropyrrol-2,5-diones with  $\alpha$ -furylmethanol. *Butlerov Communications*. **2017**. Vol.51. No.8. P.71-75. DOI: 10.37952/ROI-jbc-01/17-51-8-71