

The nature of changes in biometric indices and the activity of oxidant enzymes of seedlings after the complex effect of physical factors and synthetic growth regulators on barley seeds of ordinary (*Hordeum vulgare*)

© Viktoria M. Posokhova,¹ Pyotr P. Purygin,^{1*} Tatiana I. Vasilyeva,² and Valery F. Putko³

¹ Department of Organic, Bioorganic and Medicinal Chemistry; ² Department of Biochemistry, Biotechnology and Bioengineering. Natural Science Institute. Samara National Research University.

Moscow Ave., 34. Samara, 443086. Samara Region. Russia.

Phone: +7 (846) 334-54-59. E-mail: puryginpp2002@mail.ru, vastaty@rambler.ru

³ Innovation Center "Ecotechnology", Leninskaya St., 168, office 234. Samara, 443041. Samara Region. Russia. E-mail: vfputko@yandex.ru

*Supervising author; ⁺Corresponding author

Keywords: complex of physical factors, yield, pre-treatment, magnetic field, peroxidase, catalase, oxidant enzymes.

Abstract

This article is directed to a research of influence of preseeding processing of seeds by a complex of physical and chemical methods on biological and biochemical parameters of sprouts of barley of ordinary (*Hordeum vulgare*). The purpose of preseeding processing – release of sowing material from causative agents of diseases, increase in viability of seeds and acceleration of their germination. Long-term theoretical and pilot studies showed that the nucleus can be brought out of a condition of biological rest, influencing seeds before crops by factors, various by the nature.

In the present article we investigated and studied complex influence chemical (a humate To, a growth factor of Epina-estr and Amylum) and physical (an ultra-violet radiation, ozone and the constant magnetic field (MF) with an induction of 300 Gs) factors of preseeding processing on body height and biochemical indicators of sprouts of barley ordinary.

Seeds of barley divided into groups and irradiated in the magnetic and plasma MPU-1 installation. Then, these groups in turn processed chemical components, couched and observed for thirteen days.

Influence of these influences of factors on activity of such plants of enzymes, important for activity, as a catalase and peroxidase in leaves of sprouts of barley ordinary is studied.

The activity of peroxidase was determined by a colorimetric method by Boyarkin.

The specific activity of a catalase was determined by Korolyuk and Ivanova's method.

Preseeding processing, including a magnetic field with an induction of 300 Gs within 1 minute turned out what the increased viability of seeds, increase in length of leaves twice, activities of a catalase for 12%, peroxidases for 23% in relation to monitoring testifies to a complex of physical factors of the favorable for sprouts of barley.

The most efficient way of processing of plants can be put into practice in the modern crop production for increase in productivity of crops and increase in effectiveness of the applied mineral fertilizers.

Results of work make a particular contribution to development and perfecting of methods of preseeding processing of seeds that will allow to increase efficiency of a harvest of crops.

References

- [1] E.V. Nikolaev, V.D. Pismenny, A.V. Ryumshin. Grain Processing in Agricultural Enterprises. Edited by E.V. Nikolaev. *Simferopol*. **2008**. 209p. (russian)
- [2] N.N. Tretyakov, E.I. Koshkin, N.M. Makrushin and others. Physiology and biochemistry of agricultural plants. Ed. N.N. Tretyakov. *Moscow: Kolos*. **2000**. 34p. (russian)
- [3] V.F. Putko, A.I. Isakov, A.N. Kalimullin. Device for presowing treatment of seeds. *Russian Patent* 5037307/15, **1994**. *Bull.* No.1. 01/10/96. (russian)

- [4] P.P. Purygin, D.A. Tsaplev, V.A. Purygin, Yu.P. Zarubin, and T.I. Vasil'yeva. The study of the level of carotenoids, chlorophyll *a* and *b* in seedlings of common barley (*Hordeum vulgare*) after treatment of seeds constant magnetic field and UV radiation in the presence of ozone. *Butlerov Communications*. **2015**. Vol.42. No.5. P.23-25. ROI: jbc-02/15-42-5-23
- [5] P.P. Purygin, D.A. Tsaplev, E.V. Tsapleva, and Yu.P. Zarubin. Determination of the specific activity of peroxidase of common barley (*Hordeum vulgare*) and common millet (*Panicum miliaceum*) when exposed to ozone and constant magnetic field. *Butlerov Communications*. **2013**. Vol.35. No.9. P.90-93. ROI: jbc-02/13-35-9-90
- [6] N.A. Klyonova, O.N. Makurina, E.V. Pisareva, M.Yu. Yazykova. Special practical work on the biochemistry of animals, plants and microorganisms. *Samara: Ed. "Sprint"*. **2013**. 147p. (russian)
- [7] I.Yu. Sakharov. Palm peroxidase. *Biochemistry*. **2004**. Vol.69. No.8. P.1013-1020. (russian)
- [8] T.V. Chirkova. Physiological basis of plant resistance. *SPb .: Publishing house of S.-Petersburg Univ.* **2002**. 244p. (russian)
- [9] A.M. Avalbaev, R.A. Yuldashev, F.M. Shakirova. The physiological effect of phytohormones class brassinosteroids on plants. Successes of modern biology. Ufa. Inst. Biochemistry and genetics Ufa scientific. *Center of Sciences*. **2006**. Vol.126. No.2. P.192-200. (russian)