

## Influence of carbonate salinity on some indicators of water exchange of the triticale of the winter variety "Tribune"

© Tatiana N. Evgrashkina,<sup>1</sup> Viktor V. Ivanishchev,<sup>1\*</sup>  
Olga I. Boikova,<sup>2+</sup> and Nikolay N. Zhukov<sup>1</sup>

<sup>1</sup> Department of Biology and Technologies of Living Systems. Tula State Lev Tolstoy Pedagogical University. Lenin Ave., 125. Tula, 300026. Russia. Phone: +7 (953) 957-63-06. E-mail: [tatyanale9339@gmail.com](mailto:tatyanale9339@gmail.com)

<sup>2</sup> Department of Chemistry. Tula State Lev Tolstoy Pedagogical University. Lenin Ave., 125. Tula, 300026. Russia. Phone: +7 (906) 624-32-46. E-mail: [benosi@mail.ru](mailto:benosi@mail.ru)

\*Supervising author; +Corresponding author

**Keywords:** triticale winter (x *Triticosecale*), hydrogen peroxide, malonic dialdehyde, antioxidant protection enzymes.

### Abstract

Effect of salt stress induced by carbonate salinity caused by the presence of Na<sub>2</sub>CO<sub>3</sub> in a concentration of 120 mM was studied. As a research object were taken Triticale of the winter variety "Tribune". All measurements were carried out on 7-day triticale sprouts with exposure for 12, 24, 48, 72, 96 hours. The parameters of osmotic potential, moisture content, and sprouting length were studied. Carbonate salinity caused changes in all parameters. Thus, during the experiment, almost a twofold decrease in the osmotic potential in triticale shoots was noted for 96 hours of exposure, while in the roots this index changed insignificantly, remaining at the level of control values. At 72 hours exposure, the osmotic potential in the roots was increased by 30%. At the same time, during the first day of salinity, the moisture content in shoots was reduced by 25%, and in roots by 11%. Subsequently, by 96 hours of salinity, this index increased both in shoots and in roots, and by the end of the experiment it was 10% lower than the control one. Also, carbonate salinity adversely affected the growth of plants, causing its inhibition. This was manifested in a decrease in the length of the experimental shoots in comparison with the control samples. The growth activity of the roots was also reduced. Based on the results obtained, it can be concluded that the carbonate ions have a negative effect on winter triticale plants under carbonate salinity. This may be due to partial hydrolysis of sodium carbonate and alkalization of the medium, which is an additional stress factor for plants.

### References

- [1] A.O. Shelkova, L.P. Stepanova. Agroecological assessment of the formation of solontsi, their composition and properties. *Agrobusiness and ecology*. **2015**. No.2. P.223-225. (russian)
- [2] N.S. Marechal Patil, V.T. Apradh, B.A. Karadge. Effects of alkali stress on seed germination and seedlings growth of *Vigna Conitifolia* (Jacq.). *Article in Pharmacognosy Journal*. **2013**. No.4. P.73-76.
- [3] J. Anim. Physiological adaptive mechanisms of *Leymus chinensis* during germination and early seedling stages under saline and alkaline conditions. **2014**. *Plant Sci*. Vol.24. No.3. P.241-244.
- [4] Z. Biea, T. Itob, Y. Shinoharaba. Effects of sodium sulfate and sodium bicarbonate on the growth, gas exchange and mineral composition of lettuce *Scientia Horticulturae*. **2004**. No.99. P.215-224.
- [5] P. Warne, R.D. Guy, L. Rollins, and D.M. Reid. The effects of sodium sulphate and sodium chloride on growth, morphology, photosynthesis, and water use efficiency of *Chenopodium rubrum*. *Can J. Bot*. **1998**. No.68. P.999.
- [6] S. Renault, C. Croser, J.A. Franklin. Effects of NaCl and Na<sub>2</sub>SO<sub>4</sub> on red-osier dogwood (*Cornus stolonifera* Michx) seedlings. *Plant and Soil*. **2001**. No.233. P.261-268.
- [7] Tattini, R. Gucci, M.A. Coradeschi, C. Ponzio and J.D. Everard. Growth, gas exchange and ion content in Oleaeuropaea plants during salinity stress and subsequent relief Massimiliano. *Physiologia Plantarum*. **1995**. No.95. P.203-210.
- [8] T.N. Evgrashkina, V.V. Ivanishchev, O.I. Boikova, and N.N. Zhukov. Manifestations of oxidative stress in seedlings of triticale under conditions of sulfate salinity. *Butlerov Communications*. **2018**. Vol.54. No.4. P.128-133. ROI: jbc-02/18-54-4-128

- [9] A.R. Garifzyanov, N.N. Zhukov, V.V. Ivanishchev, A.A. Kozobryuhov. Regulation water metabolism in triticosecale in NaCl-salinity. *Visnik Kharkiv National Agrarian University. Gray biology*. **2013**. Iss.1(28). P.34-43.
- [10] N.M. Yurtaeva. Small plant physiology practicum: study manual for universities; Nizhegor. state architectures. build un-t *N. Novgorod: NNGASU*. **2015**. 112p. (russian)
- [11] A.M. Mohamed, G.N. Raldugina, V.P. Kholodova, V.V. Kuznetsov. Accumulation of osmolytes by plants of different genotypes of rapeseed during chloride salinity. *Plant physiology*. **2006**. Vol.53. No.5. P.732-739. (russian)
- [12] Yu.V. Orlova, N.A. Myasoedov, Yu.V. Balnokin. The role Na +, K +, Cl - ions and sugars in the regulation of osmotic pressure in Artemisia plants during chloride salinity. *VI Congress OFR, International Conference "Plant Physiology of the 21st Century: from molecules to ecosystems."* Syktyvkar. **2007**. Vol.2. P.307-308. (russian)
- [13] N.N. Zhukov. Study of physiological and biochemical mechanisms of salt stress in triticale in the early stages of ontogenesis. *Author. Diss ... Cand. biol. sciences. Pushchino*. **2013**. 22p. (russian)
- [14] A.A. Belozerova, N.A. Bome. Study of spring wheat reaction to salinity on the variability of sprouts morphometric parameters. *Fundamental research*. **2014**. No.12. P.300-306. (russian)
- [15] S. Keshavkant, J. Padhan, S. Parkhey, S. C. Naithani. Physiological and Antioxidant Responses of Germinating Cicer arietinum Seeds to Salt Stress. *Plant physiology*. **2012**. Vol.59. No.2. P.232-237.
- [16] D. Khasan, I.S. Kovtun, M.V. Efimova. The effect of chloride salinity on seed germination and growth of seedlings of Brassica napus L. *Vestnik Tomsk State University*. **2011**. No.4. P.108-113. (russian)
- [17] F. Yasar, S. Elialtiglu, K. Ildis. The effect of salinity on antioxidant defense systems, lipid peroxidation and chlorophyll content in the leaves of beans. *Plant physiology*. **2008**. Vol.55. No.6. P.869-874. (russian)