

## Research of the sorption activity of copper phthalocyanine

© Andrey A. Degtyarev,<sup>1\*</sup> Alexandra V. Trishina,<sup>2</sup> and Alexander G. Tarakanov<sup>3</sup>

Department of Chemistry and Chemical Technologies. Tambov State Technical University. Soviet St., 106.

Tambov, 392000. Russia. Phone: <sup>1)</sup> (960) 660-82-93; <sup>2)</sup> (920) 484-31-85; <sup>3)</sup> (953) 717-46-83.

E-mail: <sup>1)</sup> ad.dycost@gmail.com; <sup>2)</sup> koroleva\_tambov@mail.ru; <sup>3)</sup> uazqaaz@gmail.com

\*Supervising author; <sup>+</sup>Corresponding author

**Keywords:** adsorption, active centers, DFT, phthalocyanine cuprum, molecular electrostatic potential, theory of boundary orbitals, dipole moment.

### Abstract

In the work a research the sorption capacity of copper phthalocyanine with respect to polars sorbats by the method of quantum-chemical modeling. The theory of the density functional with the hybrid functional B3LYP5 and basic sets 6-31G(d,p) (optimization of geometry) and cc-pVDZ (single point calculation) was chosen as the calculation method. The purpose of this work was theoretical prediction of the effectiveness of surfactants in the hydrophobization of the copper phthalocyanine surface (necessary for the flushing process).

Modeling of sorption activity was carried out on a single molecule of copper phthalocyanine and bimolecular complex with a geometry of the crystal grid of the  $\beta$ -modification.

Prediction of sorption activity was carried out by the methods of dipole moments, molecular electrostatic potential and the frontier molecular orbital theory.

As a result of modeling, the electrophilic characteristics of the entire surface of copper phthalocyanine were established, but the sorption of nucleophilic molecules is possible only on surfaces containing indole rings. On other surfaces possible sorption of small electrophilic particles, such as OH<sup>-</sup>.

The most probable centers of sorption are the nitrogen atoms of the indole ring (confirmed by all three methods).

To the results of the research, the most effective surfactants for hydrophobizing the surface of copper phthalocyanine are nucleophilic molecules that give an acidic or neutral medium.

### References

- [1] L.M. Bykova, A.V. Eltsov, I.Y. Kvytko, L.P. Kovzhina, V.V. Shaburov, T.G. Shavva. Laboratory workshop on the synthesis of intermediate products and dyes. *Leningrad: Chemistry*. **1985**. P.352. (russian)
- [2] E.A. Indeykin, L.N. Leibzon, I.A. Tolmacheva. Pigmenting of paints and varnishes. *Leningrad: Chemistry*. **1986**. P.160. (russian)
- [3] S.N. Tolstay, S.A. Shabanova. The use of surfactants in the paint and varnish industry. *Moscow: Chemistry*. **1976**. P.176. (russian)
- [4] C. Defeyt, P.Vandenabeele, B. Gilbert, J. Van Pevenage, R. Cloots, D. Strivay Contribution to the identification of  $\alpha$ -,  $\beta$ - and  $\epsilon$ -copper phthalocyanine blue pigments in modern artists' paints by X-ray powder diffraction, attenuated total reflectance micro-fourier transform infrared spectroscopy and micro-Raman spectroscopy. *J. Raman Spectrosc.* **2012**. Vol.43. P.1772-1780.
- [5] T. Zou, X. Wang, H. Ju, L. Zhao, T. Guo, W. Wu and H. Wang Controllable Molecular Packing Motif and Overlap Type in Organic Nanomaterials for Advanced Optical Properties. *Crystals*. **2018**. Vol.8. No.22.
- [6] R. Lambourne. Paintwork materials and coatings. Theory and Practice. *St. Petersburg: Chemistry*. **1991**. 512p.
- [7] P. Kaur, R. Sachdeva, S. Singh and G. S. S. Saini DFT study of the effect of substitution on the molecular structure of copper phthalocyanine. *Citation: AIP Conference Proceedings*. **2016**. Vol.1728, 020281.
- [8] H. Vázquez, P. Jelínek, M. Brandbyge, A.P. Jauho, F. Flores Corrections to the density-functional theory electronic spectrum: copper phthalocyanine. *Appl Phys A*. **2009**. Vol.95. P.257-263.
- [9] T.V. Basova, V.G. Kiselev, Britt-Elfriede Schuster, H. Peisert and T. Chasse. Experimental and theoretical investigation of vibrational spectra of copper phthalocyanine: polarized single-crystal Raman spectra, isotope effect and DFT calculations. *J. Raman Spectrosc.* **2009**. Vol.40. P.2080-2087.
- [10] N. Marom, X. Ren, J.E. Moussa, J.R. Chelikowsky and L.K. Electronic. Structure of copper phthalocyanine from  $G_0W_0$  calculations. *PHYSICAL REVIEW*. **2011**. Vol.84. 195143p.

- [11] D. Stradi, C. Díaz, M. Alcami. A density functional theory study of the manganese-phthalocyanine. *Theor Chem Acc.* **2011**. Vol.128. P.497-503.
- [12] J.R. De Lile and S. Zhou. A density functional theory investigation of charge mobility in titanyl-phthalocyanines and their tailored peripherally substituted complexes. Препринты arXiv: 1610.10027.
- [13] A.A. Doroshenko, I.V. Nechaev and A.V. Vvedensky. Quantum-chemical modeling of the hydroxide ion gas-phase adsorption on IB metal clusters Men (n = 2-8). *Butlerov Communications.* **2015**. Vol.41. No.2. P.135-140. DOI: 10.37952/ROI-jbc-01/15-41-2-135
- [14] A.A. Degtyarev, A.G. Tarakanov. Comparison of methods for the study of water sorption on the titanium oxide. *Butlerov Communications.* **2017**. Vol.50. No.4. P.105-111. DOI: 10.37952/ROI-jbc-01/17-50-4-105
- [15] I.S. Puzyrev, E.P. Sobina, and S.V. Medvedevskih. Template synthesis and sorption of water vapor by porous silica gels with a high specific surface area. *Butlerov Communications.* **2013**. Vol.36. No.10. P.141-145. ROI: jbc-02/13-36-10-141
- [16] Yu.B. Grunin, L.Y. Grunin, E.A. Nikolskaya, and D.S. Masas. The possibility of proton magnetic relaxation in the analysis of the Gibbs adsorption of water on plant polymers. *Butlerov Communications.* **2014**. Vol.39. No.9. P.39-44. ROI: jbc-02/14-39-9-39
- [17] C. Brown Crystal Structure of P-Copper Phthalocyanine. *J. J. Chem. Soc. A.* **1968**. Vol.2488.
- [18] C. Lee, W. Yang and R.G. Parr. Phys Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Rev.* **1988**. Vol.37. P.785-789.
- [19] W.J. Hehre, R. Ditchfield and J.A. Pople Self – Consistent Molecular Orbital Methods. XII. Further Extensions of Gaussian – Type Basis Sets for Use in Molecular Orbital Studies of Organic Molecules. *J. Chem. Phys.* **1972**. Vol.56. 2257p.
- [20] V. Rassolov, J.A. Pople, M. Ratner and T.L. Windus 6-31G basis set for atoms K through Zn. *J. Chem. Phys.* **1998**. Vol.109. 1223p.
- [21] H. Thom, Jr. Dunning. Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen. *J. Chem. Phys.* **1989**. Vol.90. 1007p.
- [22] N.B. Balabanov and K.A. Peterson Systematically convergent basis sets for transition metals. I. All-electron correlation consistent basis sets for the 3d elements Sc–Zn. *J. Chem. Phys.* **2005**. Vol.123. 064107p.
- [23] Alex A. Granovsky. Firefly version 8 [Electronic resource]: Introduction to the Firefly. URL: <http://classic.chem.msu.su/gran/gamess/index.html> (access date: 16.04.2018).
- [24] E.V. Esipova Adsorption desulfurization of diesel gas condensate fuel. *Moscow.* **2015**. P.192.
- [25] L.Yu. Alikberova, E.V. Savinkina, M.N. Davydova Fundamentals of the structure of matter. Toolkit. *Moscow: MITHT.* **2004**. (russian)
- [26] R.F.W. Bader. Atoms in Molecules - A Quantum Theory. *Oxford University Press.* **1990**.
- [27] T. A. Keith AIMAll (Version 17.11.14),TK Gristmill Software [Electronic resource]: URL: [aim.tkgristmill.com](http://aim.tkgristmill.com) (access date: 20.04.2018).
- [28] N.L. Medyankin, L.A. Bodyan, I.A. Varlamova, Kh.Ya. Gireeva, N.L. Kalugina, T.A. Gireva. Study of the sorption activity of the coal surface. *Bulletin of MSTU. G.I. Nosov.* **2005**. No.3. P.11-16.
- [29] A. Pullman and B. Pullman. Molecular electrostatic potential of the nucleic acids. *Quarterly Reviews of Biophysics.* **1981**. Vol.14. No.3. P.289-380.
- [30] B.M. Bode and M.S. Gordon. *J. Mol. Graphics Mod.* **1998**. Vol.16. P.133-138.
- [31] Y. Li, Y. Liu, F. Li. Synthesis, crystal structure, vibration spectral and DFT studies of 4-aminoantipyrine and its derivatives. *Molecules.* **2013**. Vol.18. P.877-893.
- [32] H. Vázquez, P. Jelínek, M. Brandbyge, A.P. Jauho, F. Flores Corrections to the density-functional theory electronic spectrum: copper phthalocyanine. *Appl Phys A.* **2009**. Vol.95. P.257-263.
- [33] E.A. Batanova. Oxidative destruction of phthalocyanine complexes of metals in aqueous alkaline media. *Moscow.* **2001**. P.177.
- [34] H. Lee. Competitive adsorption of CO<sub>2</sub> and H<sub>2</sub>O molecules on the BaO (100) surface: A FirstPrinciple study. *Bull. Korean Chem. Soc.* **2011**. Vol.32. No.3. P.988-992.
- [35] P.C. do Couto, S.G. Estácio and B. J. Costa Cabral. The Kohn-Sham density of states and band gap of water: From small clusters to liquid water. *J. Chem. Phys.* **2005**. Vol.123. 054510p.
- [36] C.Y. Ng, D.J.Trevor, P.W. Tiedemann, S.T. Ceyer, P.L. Kronebusch, B.H. Mahan and Y.T. Lee. Photoionization of dimeric polyatomic molecules: Proton affinities of H<sub>2</sub>O and HF. *J. Chem. Phys.* **1977**. Vol.67. 4235p.
- [37] V.V. Voevodin, S.A. Zhumaty, S.I. Sobolev, A.S. Antonov, P.A. Bryzgalov, D.A. Nikitenko, K.S. Stefanov, V.V. Voevodin The practice of the supercomputer "Lomonosov". *Open systems. Moscow: Publishing house Open Systems."* **2012**. No.7. P.36-39.