

The possibilities of simple acid extraction for the extraction of certain trace elements from certified samples of coal ash and their concentrations determined by the method of optical emission spectrometry with inductively coupled plasma

Roman P. Kolmykov,^{*†} Vyacheslav E. Tsvetkov, and Anna V. Petrushina

Federal Research Center of Coal and Coal Chemistry of the SB RAS.

Sovietsky St., 18. Kemerovo, 650000. Phone: +7 (384) 228-17-21.

E-mail: kolmykoff.roman@yandex.ru

*Supervising author; †Corresponding author

Keywords: ash, ICP OES, microelements, acid extraction.

Abstract

Mineral processing with maximum efficiency is one of the most actual problems of modern economy. To solve this problem, it is necessary to develop a set of methods, including, among other methods, the analysis of the waste chemical composition. The ashes from coal combustion are promising for processing waste, since it contains a large amount of valuable trace elements.

Experimental results on the investigation of the possibility of the determination of eight micro impurity elements Co, Cr, Cu, Ni, V, Zn, La, Y in a coal ash by acid extraction using optical-emission spectrometry with inductively coupled plasma are obtained. The study provides calibration graphs for the determination of the listed elements with their concentrations in solution up to 2 mg/L, the results of determining the stability of the calibration characteristics, the determination of spectral background influences for selected analytic wavelengths in the analysis of ashes (real objects). As a result of the study, the extraction rates of these elements for the certified ash samples were determined, and the contents of the identified elements for the CO-1 ash sample missing in its passport were predicted.

When extracting the CO-1, the certain concentrations of the elements are 35–36% of their certified concentrations. In the extracts of the ZUK-1, the detectable elements are contained in amounts of from 26.7 to 41.4%. Lanthanum is poorly extracted by aqua regia, as in the case of the ZUK-2, only 1%, possibly due to the fact that it is bound to some stable chemical compound with silicon. When the ZUK-2 was extracted, the detectable elements passed into a solution in amounts ranging from 23.5 to 46.7%.

References

- [1] X. Li, S. Dai, W. Zhang, T. Li, X. Zheng, W. Chen. Determination of As and Se in coal and coal combustion products using closed vessel microwave digestion and collision/reaction cell technology (CCT) of inductively coupled plasma mass spectrometry (ICP-MS). *International Journal of Coal Geology*. **2014**. Vol.124. P.1-4.
- [2] A.K. Das, R. Chakraborty, de la M. Guardia, M.L. Cervera, D. Goswami. ICP-MS multielement determination in fly ash after microwave-assisted digestion of samples. *Talanta*. **2001**. Vol.54. No.5. P.975-981.
- [3] D.A. Spears. The use of laser ablation inductively coupled plasma-mass spectrometry (LA ICP-MS) for the analysis of fly ash. *Fuel*. **2004**. Vol.83. No.13. P.1765-1770.
- [4] K. Suzuki. Characterisation of airborne particulates and associated trace metals deposited on tree bark by ICP-OES, ICP-MS, SEM-EDX and laser ablation ICP-MS. *Atmospheric Environment*. **2006**. Vol.40. No.14. P.2626-2634.
- [5] F.G. Antes, F.A. Duarte, M.F. Mesko, M.A. Nunes, V.A. Pereira, E.I. Müller, E.M. Flores. Determination of toxic elements in coal by ICP-MS after digestion using microwave-induced combustion. *Talanta*. **2010**. Vol.83. No.2. P.364-369.
- [6] A. Smeda, W. Zyrnicki. Application of sequential extraction and the ICP-AES method for study of the partitioning of metals in fly ashes. *Microchemical Journal*. **2002**. Vol.72. No.1. P.9-16.

- THE POSSIBILITIES OF SIMPLE ACID EXTRACTION FOR THE EXTRACTION OF CERTAIN TRACE... ____ 134-140
- [7] A. Iwashita, T. Nakajima, H. Takanashi, A. Ohki, Y. Fujita, T. Yamashita. Determination of trace elements in coal and coal fly ash by joint-use of ICP-AES and atomic absorption spectrometry. *Talanta*. **2007**. Vol.71. No.1. P.251-257.
- [8] A. Iwashita, T. Nakajima, H. Takanashi, A. Ohki, Y. Fujita, T. Yamashita. Effect of pretreatment conditions on the determination of major and trace elements in coal fly ash using ICP-AES. *Fuel*. **2006**. Vol.85. No.2. P.257-263.
- [9] F. Low, L. Zhang. Microwave digestion for the quantification of inorganic elements in coal and coal ash using ICP-OES. *Talanta*. **2012**. Vol.101. P.346-352.
- [10] J. Wang, T. Nakazato, K. Sakanishi, O. Yamada, H. Tao, I. Saito. Microwave digestion with HNO₃/H₂O₂ mixture at high temperatures for determination of trace elements in coal by ICP-OES and ICP-MS. *Analytica chimica acta*. **2004**. Vol.514. No.1. P.115-124.
- [11] L. Ebdon, E.H. Evans. An introduction to analytical atomic spectrometry. *John Wiley & Sons*. **1998**.
- [12] I.J. Alinnor. Adsorption of heavy metal ions from aqueous solution by fly ash. *Fuel*. **2007**. Vol.86. P.853-857.
- [13] G. Schlemmer, B. Welz. Determination of heavy metals in environmental reference materials using solid sampling graphite furnace AAS. *Fresenius' Journal of Analytical Chemistry*. **1987**. Vol.328. No.4. P.405-409.
- [14] S. Sushil, V.S. Batra. Analysis of fly ash heavy metal content and disposal in three thermal power plants in India. *Fuel*. **2006**. Vol.85. No.17. P.2676-2679.
- [15] B.F. Arbogast. Analytical Methods Manual for the Mineral Resource Surveys Program, US Geological Survey. *US Department of the Interior, US Geological Survey*. **1996**.
- [16] R. Singh, D.P. Singh, N. Kumar, S.K. Bhargava, S.C. Barman. Accumulation and translocation of heavy metals in soil and plants from fly ash contaminated area. *Journal of Environmental Biology*. **2010**. Vol.31. P.421-430.
- [17] L. Vincze, A. Somogyi, J. Osan, B. Vekemans, S. Toeroek, K. Janssens, F. Adams. Quantitative trace element analysis of individual fly ash particles by means of X-ray microfluorescence. *Analytical chemistry*. **2002**. Vol.74. No.5. P.1128-1135.
- [18] O. Dogan, M. Koby. Elemental analysis of trace elements in fly ash sample of Yatağan thermal power plants using EDXRF. *Journal of Quantitative Spectroscopy and Radiative Transfer*. **2006**. Vol.101. No.1. P.146-150.
- [19] O. Dogan, Ö. Symsek, Y. Nuhoglu, M. Kopya, & M. Ertugrul. X-ray fluorescence spectrometry analysis of trace elements in fly ash samples of Kemerköy thermal power plants. *Journal of trace and microprobe techniques*. **2001**. Vol.19. No.2. P.289-295.
- [20] B.G. Kutchko, A.G. Kim. Fly ash characterization by SEM-EDS. *Fuel*. **2006**. Vol.85. No.17. P.2537-2544.