

## Atomic electrostatic potential on the reaction center and on the leaving group as a descriptor of processes of an aminolysis phenyl- and thiophenylacetates and hydrolysis of acetanilides

© Evgeny N. Krylov,<sup>1\*</sup> Marina V. Loginova,<sup>1</sup> Lyudmila V. Virzum,<sup>2+</sup> Tatiana A. Shapovalova,<sup>2</sup> and Matvey S. Gruzdev<sup>3</sup>

<sup>1</sup> Chemistry department. Ivanovo State University. Ermak St., 39. Ivanovo, 153025. Russia. Phone: +7 (4932) 37-37-03. E-mail: [enk2000S@yandex.ru](mailto:enk2000S@yandex.ru)

<sup>2</sup> Ivanovo State Agricultural Academy Named after D.K. Belyaev. Sovetskaya St., 45. Ivanovo, 153012. Russia. Phone: +7 (4932) 32-81-44. E-mail: [virzum@list.ru](mailto:virzum@list.ru)

<sup>3</sup> G.A. Krestov Institute of Solution Chemistry. Russian Academy of Sciences. Akademicheskaya St., 1. Ivanovo, 153045. Russia. Phone: +7 (4932) 33-62-59.

\*Supervising author; +Corresponding author

**Keywords:** molecular electrostatic potential, carbonyl atom of carbon, hydrolysis of acetanilides, aminolysis of acetates, nucleophilic attack.

### Abstract

Hydrolysis of acetanilides and aminolysis of phenyl- and thiophenylacetates are related reactions since all of them are processes of nucleophilic substitution on the carbon's carbonyl atom. Current views of chemical reactivity based on the DFT theory leaning upon reactivity indices that are descriptors of both the reaction center and the molecule as a whole have not been applied to the given processes before. One of such descriptors is an atomic electrostatic potential characterizing the distribution of electron density on the reaction center. The given parameter was calculated by the DFT theory M06/6-311+G\* (MeCN, SMD). There have been calculated with full optimization and without any restrictions as to the type of symmetry the structures of substituted phenylacetates XPhO-C(O)Me, acetanilides XPhNHC(O)Me, thiophenylacetates ZPhSC(=O)Me and benzylamines XPhCH<sub>2</sub>NH<sub>2</sub> (X and Z are substitutes). All relationships between the atomic electrostatic potential and the charge on the reaction center in the Hirshfeld scheme are symbasic. In all of the cases the rate is determined by the nucleophilic attack on the reaction center with the activity/selectivity relationship being observed. This suggests that the thiophenolate-anion is a better leaving group compared to the phenolate-anion and as well as that the anionic abstraction of the leaving group does not limit the reaction rate. The fact that the reaction rate is limited by the nucleophilic attack of the reagent is not inconsistent about concerted mechanism of reaction the views, since it is known that such reactions may be quite concerted though not quite synchronous.

### References

- [1] Chemical reactivity theory. A density functional view. P.K. Chattaraj, Ed. Boca Raton: *CRC Press*. **2009**. 610p.
- [2] Concepts and methods in modern theoretical chemistry. Electronic structure and reactivity. S.K. Ghosh, P.K. Chattaraj. Eds. N.-Y.: *CRC Press*. **2013**. 450p.
- [3] P. Politzer, J.S. Murray. The electrostatic potential: an overview. *Theor. Chem. Acc.* **2002**. Vol.108. P.134-142.
- [4] B. Galabov, S. Ilieva, B. Hadjieva, Y. Atanasov, Schaefer III H.F. Predicting Reactivities of Organic Molecules. Theoretical and Experimental Studies on the Aminolysis of Phenyl Acetates. *Phys. Chem.* **A2008**. Vol.112. P.6700-6707.
- [5] L.V. Kurit'syn, T.P.Kustova, A.I. Sadovnikov, N.V. Kalinina, M.V. Kluev. The kinetic of acyl transfer reactions. *Ivanovo. IvGU*. **2006**. 260p. (russian)
- [6] D. Cheshmedzhieva, S. Ilieva, B. Hadjieva, T. Trayanova. Reactivity of acetanilides in the alkaline hydrolysis reaction: theory vs. experiment. *Mol. Phys.* **2009**. Vol.107. No.8-12. P.1187-1192.
- [7] H.K. Oh, J.Y. Yang, H.W. Lee, I. Lee. Kinetics and Mechanism of the Aminolysis of Thiophenyl Acetates in Acetonitrile. *Bull. Korean Chem. Soc.* **1999**. Vol.20. No.12. P.1418-1420.

- [8] H.K. Oh, S.K. Kim, I. Lee. Nucleophilic Substitution Reactions of Thiophenyl Phenylacetate with Benzylamines in Acetonitrile. *Bull. Korean Chem. Soc.* **1999**. Vol.20. No.9. P.1017-1020.
- [9] M.S.J. Dewar. Multibond Reactions Cannot Normally Be Synchronous. *J. Am. Chem. Soc.* **1984**. Vol.106. No.1. P.209-219.
- [10] R.S. Brown. Studies in Amide Hydrolysis: The Acid, Base, and Water Reactions. In: The amide linkage: selected structural aspects in chemistry, biochemistry and materials science. C.M. Breneman, A. Greenberg, J.F. Liebman, Eds. *Wiley Int. N.-Y.* **1999**. P.85-115. 672p.
- [11] M. Valiev, E.J. Bylaska, N. Govind, K. Kowalski, T.P. Straatsma, van H.J.J. Dam, D. Wang, J. Nieplocha, E. Apra, T.L. Windus, de W.A. Jong. NWChem: a comprehensive and scalable open-source solution for large scale molecular simulations. *Comput. Phys. Commun.* **2010**. Vol.181. Iss.9. P.1477-1489.
- [12] A.V. Marenich, C.J. Cramer, D.G. Truhlar. Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions. *J. Phys. Chem. B.* **2009**. Vol.113. P.6378-6396.
- [13] B. Galabov, D. Cheshmedzhieva, S. Ilieva, B. Hadjieva. Computational Study of the Reactivity of N-Phenylacetamides in the Alkaline Hydrolysis Reaction. *J. Phys. Chem. A.* **2004**. Vol.108. No.51. P.11457-11462.
- [14] D. Cheshmedzhieva, S. Ilieva, B. Hadjieva, B. Galabov. The Mechanism of Alkaline Hydrolysis of Amides. A Comparative Computational and Experimental Study of the Hydrolysis of N-Methylacetamide, N-Methylbenzamide and Acetanilide. *J. Phys. Org. Chem.* **2009**. Vol.22. P.619.
- [15] K.T. Douglas. Elimination-addition pathways for thiol esters. *Acc. Chem. Res.* **1986**. Vol.19. P.186-192.
- [16] F.L. Hirshfeld. Bonded atom fragments for describing molecular charge densities. *Theor. Chim. Acta.* **1977**. Vol.44. No.1. P.129-138.
- [17] Reaction ability and reactions path. G. Klopman, Ed. *Moscow: Mir.* **1977**. 384p. (russian)
- [18] S.A. Akhnazarova, V.V. Kafarov. Methods of experimental optimization in chemistry and chemical technology. *Moscow: H.Sch.* **1985**. P.327. (russian)
- [19] E.N. Krylov. Dissociation of aromatic sulfonic acids in the gas and water phase: quantum-chemical calculation of pKa. *Proceedings of IvGU.* **2015**. Iss.1. P.32-39. (russian)
- [20] E.N. Krylov, M.S. Gruzdev, L.V. Virzum. Acidity of aromatic sulphonic acids in gas and water phase. *Butlerov Communications.* **2015**. Vol.42. No.6. P.117-123. ROI: jbc-02/15-42-6-117
- [21] A.I. Morar, E.N. Krylov, L.V. Virzum. Molecular electrostatic potential as a descriptor of aryl sulphonyl halogenides reaction ability. Abstr. of rep. at the X All-Russia school of young scientists. *Theoretical and experimental chemistry of liquid phase systems. (the Krestov readings).* **2015**. P.56-57. (russian)
- [22] E.N. Krylov. Molecular electrostatic potential as a descriptor of aryl sulphonyl halogenides reaction ability. *Proceedings of IvGU.* **2016**. Iss.2. P.61-70. (russian)
- [23] E.N. Krylov, L.V. Virzum. Hydrodehalogenation of aryl halogenides as a function of MEP. Mat. Of 12<sup>th</sup> Internat. Science-practical conference "National science in time of changes: postulates of the past and new age theory". *National association of scientists. Ekaterinburg.* **2015**. No.7(12). Part 3. P.70-73. (russian)
- [24] M.S. Gruzdev, L.V. Virzum, E.N. Krylov. Reaction of aromatic hydrodechlorination: quantum-chemical diagnostics of the mechanism. *Butlerov Communications.* **2015**. Vol.41. No.2. P.115-120. ROI: jbc-02/15-41-2-115