

Creation of lipid shells microcapsules of insecticides based on chlorpyrifos of trade mark GET

© Andrey V. Tretyakov,^{1,2+} Vyacheslav F. Markov,^{1,3*} and Sergey B. Yaroshuk²

¹Physical and Colloid Chemistry Department. Chemistry and Technology Institute. Ural Federal University named after the first President of Russia B.N. Yeltsin. Mira St., 19. Yekaterinburg, 620002. Sverdlovsk Region. Russia. Phone: +7 (343) 374-39-05.

²Limited liability company «GET Biotech». Vilonova St. 45E, 508 Yekaterinburg, 620062, p/b 31. Sverdlovsk Region. Russia. Phone: +7 (343) 328-54-38.

³Ural State Fire Service Institute of Emergency Ministry of Russia. Mira St., 22. Yekaterinburg, 620022. Sverdlovsk Region. Russia. Phone: +7 (343) 360-80-65.
E-mail: tretyakoff_93@mail.ru

*Supervising author; +Corresponding author

Keywords: insecticides, microcapsulation, liposomic vesicles, lipides.

Abstract

In this study, a laboratory synthesis of microencapsulated insecticides based on the active substance chlorpyrifos was performed, where DV was encapsulated in lipid monolamellar shells by a physico-chemical method in the pH range of 5-7. The resulting lipid vesicles showed high stability in the concentrate, the absence of agglomeration of particles of the nanoscale, and resistance to oxidation. Also, a surfactant, a high molecular weight polyvinylpyrrolidone, used to create a primary emulsion, was selected experimentally.

The composition of the shell includes phospholipids, the solvent of the active substance is ethyl alcohol, water acts as a precipitant. This method is easy to use, does not require complicated equipment and high energy costs.

The sizes of the obtained microcapsules were in the range from 50 nanometers to several micrometers. When analyzing particle sizes, there was no tendency for agglomeration, which confirms the presence of solid shells of vesicles.

Stability analysis was carried out by centrifugation, the size analysis of the microcapsules was carried out using a Nanofox device. Also, the dependence of the size of the final microcapsules on the synthesis temperature was investigated and microphotographs of the insecticide samples were made with an increase of 1500x magnification with the microscope *Micromed 2 var. 2-20*.

References

- [1] V.A. Nabokov. Contact insecticides, their properties and use in medical pest control. *Moscow: Medgis. 1958.* 258p. (russian)
- [2] O. Koul, S. Walia. Comparing impacts of plant extracts and pure allelochemicals and implications for pest control. *CAB Reviews. 2009.* Vol.4. P.1-30.
- [3] S.O. Duke, C.L. Cantrell, K.M. Meepagala, D.E. Wedge, N. Tabanca and K.K. Schrader. Natural Toxins for Use in Pest Management. *Toxins. 2010.* Vol.2. P.1943-1962.
- [4] C. Wang, M.E. Scharf, G.W. Bennett. Behavioral and physiological resistance of the German cockroach to gel baits (Blattodea: Blattellidae). *Econ Entomology. 2004.* Vol.97. No.6. P.2067-2072.
- [5] S.K. Prabhakaran, S.T. Kamble. Purification and characterization of an esterase isozyme from insecticide resistant and susceptible strains of german cockroach, *Blattella germanica* (L.). *Insect Biochemistry and Molecular Biology. 1995.* Vol.25. No.4. P.519-524.
- [6] Chlorpyrifos [Electronic resource] Access mode: http://www.pesticidy.ru/active_substance/chlorpyrifos (The date of application is 5.05.2018).
- [7] E.F. Stepanova, M.E. Kim, K.B. Murzagulova, S.B. Evseeva. Microcapsules: perspectives of use in modern pharmaceutical practice. *Modern problems of science and education. 2014.* No.5. P.22.
- [8] E.A. Petrova, A.V. Krikova, E.F. Stepanova. The study of the parameters of microencapsulation in the preparation of the prolonged form of naltrexone. *Chemical Pharmacy. 2014.* Vol.31. P.50-53.
- [9] A.S. Minina, A.A. Selicheva, G.M. Sorokoumova and others. Inclusion of rifampicin in multilayer and single-layer vesicles (liposomes) of different lipid composition. *Biophysics. 2004.* Vol.49. P.674-679.
- [10] M.A. Kiselev et al. Model of separated form-factors for unilamellar vesicles. *Application. Physics. 2002.* Vol.74. P.1654-1656.