To the conclusion of the equations of state of the oxyhydrate colloid

© Yury I. Sukharev,^{1*+} and Boris A. Markov²

¹ Chelvabinsk State University. Street Brothers Kashirin, 129. Chelyabinsk, 454001. Chelyabinsk Region. Russia. E-mail: Yury Sucharev@mail.ru ² Chelyabinsk Higher Military Aviation Red Banner School of Navigators, a Branch of Military Training Scientific Center Air Force "Professor N.E. Zhukovsky and Y.A. Gagarin Air Force Academy". Town, 11, 1. Branch of the VUNTS VVS "VVA". Chelyabinsk, 454015. Russia.

*Supervising author; ⁺Corresponding author

Keywords: emission-wave duality, Lagrangian of the display, operator Liesegang, oxihydrates systems, colloid clusters, spontaneous pulsation flow, diffuse electric double layer, topological continuum, dissociative-disproportionate to mechanism, the theory of Whitney, geometry of the caustic, noise state.

Abstract

The operator of Liesegang and the corresponding phase diagram, passing through the gel optical electromagnetic field have been considered. A spiral form of distribution of clusters in the gel space, resulting in a periodic change in the intensity of electromagnetic radiation transmitted through the gel is shown. The influence of pulsation noise or self-organization current in a magnetic field on the optical characteristics of tin oxy hydrate, as well as on the kinetic curves of changes in the optical density of oxyhydrate systems, the sorbing properties of *d*- and *f*-elements, and the structural organization of their colloids is established.

The paper considers also non-linear properties of the gel oxygenated systems. The discovery of the periodic properties including sorption characteristics detected by electric fluctuation conductivity on the background of the polarization phenomena.

The answer to the question about the nature of oxyhydrate gels from the position of colloidal chemical representations is received. The performed quantum chemical calculations of oxyhydrate systems indicate macromolecular binding of some single elements in the spatial gel grid. Numerous data confirm the wellknown model of the structure of co gels continuous spatial matrix that includes cells filled micellar environment.

The effect of pulsation noise or self-organization current of gel systems in a magnetic field on the optical characteristics of yttrium oxyhydrate, for example, as well as on the kinetic curves of changes in the optical density of oxyhydrate systems, as well as their structural organization, determining the properties of dand f-elements, is shown. The excited systems in the presence of noise demonstrate the effect of coherent resonance of ion-cluster flows.

An analogue of the generalized schrödinger equation for colloids is obtained. The derivation of the equation is based on an experiment that allows us to find the oscillations of the colloid with a very high degree of reliability.

The analogue of the schrödinger equation allowed us to give a holistic description of the wave, oscillatory nature of the colloidal substance. Based on this approach, it is possible to apply experimental data to restore the interaction of substances in the colloid using the mathematical apparatus of its own functions, to find interacting substances and in some way to evaluate their characteristics.

References

- [1] Passport 422272-270-42885515 PS. Conversion media multifunctional E-270. Moscow, CJSC "L-CARD". (russian)
- [2] G.I. Volovich. Circuit design analog and hardware-digital electronic devices. *Moscow: Izd. Dodek's House - XXI.* 2005. 528p. (russian)
- [3] Yu.I. Sukharev. Holographic interpretation of clusters of related water in gels of iron oxyhydrate(III), yttrium and aluminum. Butlerov Communications. 2017. Vol.52. No.10. P.28-44. DOI: 10.37952/ROI-jbc-01/17-52-10-28
- [4] Yu.I. Sukharev, and B.A. Markov. Reconstruction of spontaneous electric current attractors in oxyhydrides of *d*-elements and phonon (or electro-acoustic) echo in colloidal systems. *Butlerov* Communications. 2017. Vol.51. No.9. P.20-44. DOI: 10.37952/ROI-jbc-01/17-51-9-20

Full Paper

- [5] Yu.I. Sukharev, I.Yu. Apalikova, and O.M. Shamina. Giant clusters of distilled water in ratchet forming coxeter space. Butlerov Communications. 2015. Vol.41. No.2. P.47-55. DOI: 10.37952/ROI-jbc-01/15-41-2-47
- [6] Yu.I. Sucharev, B.A. Markov, and O.M. Shanina. New principles of the research into imperfect crystallographic forms of colloidal chemical clusters. Butlerov Communications. 2013. Vol.36. No.11. P.30-43. ROI: jbc-02/13-36-11-30
- [7] B.A. Markov, Yu.I. Sukharev, I.Yu. Apalikova. Caustics of stochastic flow clusters and a solution to the problem of formation of cluster particles themselves. Butlerov Communications. 2012. Vol.31. No.8. P.117-124. ROI: jbc-02/12-31-8-117
- [8] Yu.I. Sucharev, O.M. Krutikova, M.B. Azarov, and T.I. Prolubnikova. The spatial organization of the giant clusters of distilled and deionized natural water. Butlerov Communications. 2011. Vol.28. No.18. P.11-32. ROI: jbc-02/11-28-18-11
- [9] Yu.I. Sukharev, I.Yu. Apalikova, and O.M. Shamina. Giant clusters of distilled water in ratchet forming coxeter space. Butlerov Communications. 2015. Vol.41. No.2. P.47-55. DOI: 10.37952/ROI-jbc-01/15-41-2-47
- [10] Yuri I. Sucharev. Wave Oscillations in Colloid Oxyhydrates. Switzerland, UK, USA: Trans Tech Publications.LTD. 2010. P.497.
- [11] V.S. Anishchenko. Nonlinear effects in chaotic and stochastic systems. *Moscow: Izhevsk: In-t* computer.insled. 2003. 544p. (russian)
- [12] Application for patent No. 2018126981 "Method of research of nanocurrent ferroelectric manifestations of gels of oxyhydrates of d- and f-elements and a device for the detection of such nanocurrent pulsating ferroelectric manifestations" with priority from 07.23.2018".
- V.I. Arnold. Caustics and Wave Fronts Singularities. Moscow: Fazis. 1996. 562p. [13]
- [14] V.I. Arnold. Catastrophe Theory. Moscow: Editorial URSS. 2004. 128p. (russian)
- Yu.I. Sukharev, Ya.N. Lepp. On the sorption characteristics of oxyhydrates of some rare earth elements. [15] Neorg mater. 1995. Vol.31. No.12. P.1562-1565. (russian)
- [16] Ch. Amphlett. Inorganic Ionites. Moscow: Mir. 1966. 188p. (russian)
- Yu.I. Sukharev. Synthesis and application of specific oxyhydrate sorbents. Moscow: Energoatomizdat. [17] **1987**. 120p. (russian)
- [18] Yu.V. Yegorov. Statics of sorption of microcomponents by oxyhydrates. *Moscow: Atomizdat.* 1975. 218p. (russian)
- [19] N. Mironov. On the formation of basic salts and lanthanum hydroxide. Log neorgan. Chemistry. **1966**. Vol.XI. Iss.3. P.458. (russian)
- [20] G. Tsundel. Hydration and intermolecular interaction. *Moscow: Mir.* 1972. 310p. (russian)
- Yu.I. Sukharev, Ya.N. Lepp. On the sorption characteristics of oxyhydrates of some rare earth elements. [21] Neorg mater. 1995. Vol.31. No.12. P.1562-1565. (russian)
- Y.I. Sukharev and B.A. Markov. Liesegang rings as the common gross property of oxyhydrate and [22] other gel polymer systems: another look at the problem of periodicity. *Molecular Physics.* 2004. Vol.102. No.7. P.745-755.
- [23] B.A. Markov, Yu.I. Sukharev. Liesegang Operator. Liesegang Rings as the Common Ross-Property of Oxyhydrate Gel Polymer Systems. Известия ЧНЦ УрО РАН. 2002. №2. С.54 (www.csc.ac.ru/news).
- [24] M. Tabor. Chaos and integrability in nonlinear dynamics. *Moscow: Editorial URSS*. 2001. 320p. (russian)
- S. Chandrasekhar. Liquid crystals. Moscow: Mir. 1980. 344p. (russian) [25]
- A.S. Sonin. Introduction to the physics of liquid crystals. Moscow: Science. 1983. (russian) [26]
- Yu.I. Sukharev, B.A. Markov, I.V. Antonenko. Circular autowave pacemakers in thin-layered [27] zirconium oxyhydrate. J. Chemical Physics Letters 356 (2002) 55-62p.
- A.M. Ilyin, B.A. Markov. Nonlinear Equation of Diffusion and the Liesegang Ring. The Academy of [28] Sciences Papers. 2011. Vol.440. No.2. P.164-167.