

Thematical course: Use of biopulping for pretreatment of wood in bioethanol production. Part 1.

## Sampling of strains of basidiomycetes and micromycetes for biopulping of wood substrates

Sharapova Irina Edmundovna

Biochemistry and Biotechnology Laboratory. Biology Institute of Komi Scientific Center of Ural Department of the Russian Academy of Sciences. Kommunisticheskaya St., 28. Syktyvkar, 167982. Komi Republic. Russia.

Phone: +7 (904) 100-19-55. E-mail: scharapova@ib.komisc.ru

**Keywords:** biopulping, fungi, lignocellulosic materials, solid state fermentation, oxidase and cellulase activity.

### Abstract

Prospects of using producer strains for biopulping is evidenced by plant materials' bioconversion ability due to a complex of lignocellulolytic enzymes. The research was performed using basidiomycetes and micromycetes to determine the most promising strains for pretreatment (biopulping) of various wood substrates along with chemical treatment methods. Screening of some strains of fungi of *Trichoderma*, *Fusarium*, *Penicillium*, *Beauveria* genera has been performed to find their ability to degrade lignocellulose in accordance with testing methods. Strains with maximum oxidase and cellulase activity on agar media with gallic acid and carboxymethyl cellulose have been selected out of the group of micromycetes. In the order of oxidase activity increase, Selected strains are positioned as follows: *Fusarium sp.* < *Penicillium sp.* ≤ *Trichoderma lignorum* (F-98) < *Beauveria bassiana* (F-145). In the order of cellulase activity increase, these strains are positioned as follows: *Fusarium sp.* < *Beauveria bassiana* < *Penicillium sp.* ≤ *Trichoderma lignorum*. Comparative assessment of growth of selected strains of fungi during solid phase fermentation on softwood pulp in the course of 7, 14 and 23 days. Modification of lignocellulosic substrates with chemicals was performed using alkali and acid, as well as with organic solvents. It was found that softwood dust after treatment by alkaline hydrolysis, as well as after resin extraction and alkali treatment, is the most adapted substrate for solid state fermentation with selected strains of fungi. For the first time, use of strain of *Beauveria bassiana*, which belongs to entomopathogenic fungi group, has been considered for the purposes of biopulping. Strains of micromycetes *Penicillium sp.* and *Trichoderma lignorum*, as well as strains of basidiomycetes *Panus tigrinus* (F-8/18) and *Pleurotus eryngii* have been found that demonstrate high lignocellulosic activity and bioconversion ability of poorly hydrolysable wood substrates.

### References

- [1] R. Arora, S. Behera, S. Kumar. Bioprospecting thermophilic/thermotolerant microbes for production of lignocellulosic ethanol: a future perspective. *Renewable and Sustainable Energy Reviews*. **2015**. Vol.51. P.699-717.
- [2] Z. Xu, F. Huang. Pretreatment methods for bioethanol production. *Applied Biochemistry and Biotechnology*. **2014**. Vol.174. No.1. P.43-62.
- [3] J.N. Putro et al. Pretreatment and conversion of lignocellulose biomass into valuable chemicals. *RSC Advances*. **2016**. Vol.6. No.52. P.46834-46852.
- [4] M. Dashtban, H. Schraft, and W. Qin. Fungal bioconversion of lignocellulosic residues; opportunities & perspectives. *Int. J. Biol. Sci.* **2009**. Vol.5. P.578-595.
- [5] Lignins (structure, properties and reactions). Ed. Sarkanyan K.V., Lyudvig K.Kh. *Moscow: Forest Industry*. **1975**. 652p. (russian)
- [6] M. Madadi, Y. Tu, A. Abbas. Pretreatment of lignocellulosic biomass based on improving enzymatic hydrolysis. *Int J Appl. Sci. Biotechnol.* **2017**. Vol.5. No.1. P.1-11.
- [7] R. Sharma, V. Palled, R.R. Sharma-Shivappa, and J. Osborne. Potential of potassium hydroxyde pretreatment of switchgrass for fermentable sugar production. *Appl. Biochem. Biotechnol.* **2011**. Vol.28. P.616-626.
- [8] G. Banerjee et al. Rapid optimization of enzyme mixtures for deconstruction of diverse pretreatment / biomass feedstock combinations. *Biotechnology for Biofuels*. **2010**. Vol.3. No.1. P.22.
- [9] C.A. Rezende et al. Chemical and morphological characterization of sugarcane bagasse submitted to a delignification process for enhanced enzymatic digestibility. *Biotechnology for Biofuels*. **2011**. Vol.4. No.1. P.54.

- SAMPLING OF STRAINS OF BASIDIOMYCETES AND MICROMYCETES FOR BIOPULPING...* \_\_\_\_\_ 140-145
- [10] A.S. Meyer, L. Rosgaard, H.R. Sørensen. The minimal enzyme cocktail concept for biomass processing. *Journal of Cereal Science*. **2009**. Vol.50. No.3. P.337-344.
- [11] S. Behera, R. Arora, S. Kumar. Bioprospecting the cellulases and xylanases thermozyms for the production of biofuels. In: Paper presented at AIChE annual meeting, *San Francisco*. **2013**. 3-8 November.
- [12] L.Q. Guo, S.X. Lin, X.B. Zheng et al. Production, purification and characterization of a thermostable laccase from a tropical white-rot fungus. *World J. Microbiol Biotechnol*. **2011**. Vol.27. P.731-735.
- [13] P. Giardina et al. Laccases: a never-ending story. *Cellular and Molecular Life Sciences*. **2010**. Vol.67. No.3. P.369-385.
- [14] X. Xu, M. Lin, Q. Zang, S. Shi. Solid state bioconversion of lignocellulosic residues by *Inonotus obliquus* for production of cellulolytic enzymes and saccharification. *J. Bioresource Technology*. **2018**. Vol.247. P.88-995.
- [15] A.V. Bolobova, A.A. Askadsky, V.I. Kondratenko, M.L. Rabinovich. Theoretical bases of woody composites biotechnology. V. 2. Ferments, models, processes. *Moscow: Science Publicity*. **2002**. (russian)
- [16] T.V. Rusinova. Development of the biosynthesis technology of laccase by basidiomycetes from the *Trametes* genus. *The Author's Paper...Candidate of Technical Sciences*. *Moscow*. **2007**. (russian)
- [17] M.L. Rabinovich, A.V. Bolobova, L.G. Vasilchenko. Decomposition of natural aromatic structures and xenobiotics with fungi (review). *Applied Biochemistry and Microbiology*. **2004**. Vol.40. No.1. P.5-23. (russian)
- [18] N.S. Mokrushina, T.S. Tarasova, I.V. Darmov. Bioconversion of wood waste by composting with production of organic fertilizer. *Proceedings of the Samara Science Centre RAS*. **2009**. Vol.11. No.1. P.228-232. (russian)
- [19] *Patent RF 2355753*. The strain of *Penicillium sizovae* as a wood lignin destructor. Mokrushina N.S., Darmov I.V. No.2007126013/13; applied 09.07.2007; published 20.05.2009. *Bul.* No.14. (russian)
- [20] I.S. Geles. Wood raw as a strategic basis and reserve of civilization: *Petrozavodsk: Karelsky Science Centre RAS*. **2007**. 499p. (russian)
- [21] D.A. Satton, A.V. Fotergill, M.G. Rinaldi. Key book for pathogenic and conditionally pathogenic fungi: Trans. from English. *Moscow: World Publicity*. **2001**. 486p. (russian)
- [22] M.A. Litvinov. Key book for microscopic soil fungi. *Moscow: Science Publicity*. **1969**. 303p. (russian)
- [23] R.M. Teather, P.J. Wood. Use of congo-red polysaccharide interaction in enumeration and characterization of cellulolytic bacteria the bovine rumen. *Appl. Environ Microbiol*. **1982**. Vol.43. P.777-780.
- [24] I.V. Solovyeva, O.N. Okunev, V.V. Velkov et al. Production and properties of *Penicillium verruculosum* mutants – superproducers of cellulases and xylanases. *Microbiology*. **2005**. Vol.74. No.2. P.172-178. (russian)
- [25] *Methods of experimental mycology*. Ed. V.I. Bilai. *Kiev: Naukova dumka*. **1982**. 552p.
- [26] N.B. Gradova et al. Laboratory Practicum on General Microbiology. *Moscow: D.I. Mendeleev Russian University of Chemistry and Technology*. **1999**. 130p. (russian)
- [27] A.V. Obolenskaya et al. Practical works on wood and cellulose chemistry. *Moscow: Forest Industry*. **1965**. 320p. (russian)