

Effect of 1,2,4-triazole derivatives structure on their sorption by hypercarb under RP HPLC conditions

© Yulia P. Kurnysheva, Semen A. Ryzhkin and Svetlana V. Kurbatova*[†]

Department of Physical Chemistry and Chromatography, Samara National Research University.

Acad. Pavlova St., 1. Samara, 443011. Russia. Fax: +7 (846) 334-54-17. E-mail: curbatsv@gmail.com

*Supervising author; [†]Corresponding author

Keywords: high performance liquid chromatography, 1,2,4-triazole derivatives, porous graphitized carbon – hypercarb, aqueous acetonitrile solution, chromatographic retention, sorption thermodynamics.

Abstract

The results of 1,2,4-triazole derivatives studies by high-performance liquid chromatography using water-acetonitrile mixture in various volume ratios as eluent are presented. A porous graphitic carbon – hypercarb was used as a sorbent. It has been shown that the main physicochemical parameters determining the sorption of 1,2,4-triazole derivatives under the conditions of RP HPLC are the lipophilicity, polarizability and the volume of analyte molecules. At the same time, almost all the investigated substances are relatively polar, which contributes to their interaction with the components of the polar mobile phase. In general, among the derivatives of 1,2,4-triazole selected for the study, the values of the molecular volume, polarizability, lipophilicity and dipole moment vary within relatively wide limits, which leads to significant differences in the sorption characteristics of these compounds, which are also determined by the nature of the sorbent. At the same time, a symbiotic relationship was established in changing the values of the retention factor of the studied substances, their polarizability and lipophilicity. The observed deviations from the dependences obtained are due to specific interactions of sorbate molecules with the components of the mobile phase. The effect of temperature on the chromatographic retention of azoles was investigated. The sorption isotherm of 2-(1H-triazol-1-ylmethyl) phenol is obtained, which is almost linear in the initial part, which corresponds to the Henry region. The thermodynamic sorption characteristics of triazole derivatives have been experimentally determined. It is established that the values of changes in standard molar enthalpy and Gibbs energy of sorption during the transition of the molecules of the analyzed analytes from the bulk solution to the surface layer of the hypercarb are predominantly negative, which indicates the exothermic process and its spontaneous flow. At the same time, these values lie in a relatively narrow range and are determined mainly by the nature of the substituent and its position in the heterocycle.

References

- [1] A. Wang, P.W. Carr. Comparative study of the linear solvation energy relationship, linear solvent strength theory, and typical-conditions model for retention prediction in reversed-phase liquid chromatography. *J. Chromatogr. A* **2002**. Vol.965. P.3-23.
- [2] M.H. Abraham, A. Ibrahim, A.M. Zissimos. Determination of sets of solute descriptors from chromatographic measurements. *J. Chromatogr. A* **2004**. Vol.1037. P.29-47.
- [3] V.D. Shatz, O.V. Sahartova. High performance liquid chromatography. *Riga: Zinatne*. **1988**. 390p.
- [4] R. Berges, V. Sanz-Nebot, J. Barbosa. Modelling retention in liquid chromatography as function of solvent composition and pH of the mobile phase. *J. Chromatogr. A*. **2000**. Vol.869. No.1-2. P.27-39.
- [5] F. Gritti, G. Guiochon. Thermodynamics of adsorption of binary aqueous organic liquid mixtures on a RPLC adsorbents. *J. Chromatogr. A*. **2007**. Vol.1155. P.85-99.
- [6] B.R. Sayfutdinov, V.A. Davankov, M.M. Ilyin. Thermodynamic characteristics of the adsorption of 1,3,4-oxadiazoles and 1,2,4,5-tetrazines from methanol and water-methanol solutions on hypercrosslinked polystyrene. *Journal of Physical Chemistry*. **2017**. Vol.91. No.3. P.539-543. (russian)
- [7] S.A. Dzhabieva, S.V. Kurbatova. The effect of the nature of the sorbent on the retention of benzotriazole derivatives under RP-HPLC conditions. *Journal of Physical Chemistry*. **2015**. Vol.89. No.10. P.1644-1650. (russian)

- [8] Dzhabieva Sara Ali kyzy, E.A. Kolosova, I.N. Karaseva, M.O. Karasev, and S.V. Kurbatova. Water acetonitrile solutions features of certain aromatic heterocycles. *Butlerov Communications*. **2016**. Vol.46. No.5. P.54-60. DOI: 10.37952/ROI-jbc-01/16-46-5-54
- [9] V.I. Ivansky. Chemistry of heterocyclic compounds. *Moscow: Higher school*. **1978**. 559p. (russian)
- [10] S.N. Lanin, M.Yu. Ledenkova, Yu.S. Nikitin. Measurement of the isotherms of adsorption of anisole and benzene from a solution in n-hexane on hydroxylated silica gel by high performance liquid chromatography. *Vestnik Mosk. University. Ser. 2. Chemistry*. **1998**. Vol.39. No.5. (russian)
- [11] N.A Nekrasova, S.V. Kurbatova. Comparative Study of Quinolines and Tetrahydroquinolines Sorption on Various Sorbents from Water–Acetonitrile Solutions. *Journal of Chromatographic Science*. **2019**. Vol.57. No.4. P.369-380.
- [12] T. Hanai. Separation of polar compounds using carbon columns. *J. Chromatogr. A*. **2003**. Vol.989. P.183-196.
- [13] S. Bocian, J. Soukup, P. Jandera, B. Buszewski. Thermodynamics Study of Solvent Adsorption on Octadecyl-Modified Silica. *Chromatographia*. **2015**. Vol.78. P.21-30.